



ISSN (E): 2277- 7695  
 ISSN (P): 2349-8242  
 NAAS Rating: 5.03  
 TPI 2019; 8(5): 773-778  
 © 2019 TPI  
 www.thepharmajournal.com  
 Received: 09-03-2019  
 Accepted: 13-04-2019

**Yogeshwar Sharma**  
 Associate Professor, Department  
 of Chemistry, Motilal Nehru  
 College, Benito Juarez Marg,  
 University of Delhi South  
 Campus, New Delhi, India

## A kinetic study on the complexation of Cu (II) with $\gamma$ -aminobenzoic acid

**Yogeshwar Sharma**


### Abstract

A kinetic study on interaction of Cu(II) with  $\gamma$ -aminobenzoic acid has been investigated at 25, 30, 35 and  $40 \pm 0.05^\circ\text{C}$  in the pH range 2.01-3.14, using Aminco Morrow Stopped Flow Spectrophotometer. The ionic strength was maintained at 0.1 M  $\text{KNO}_3$ . The deprotonated form of  $\gamma$ -aminobenzoic acid is more reactive and the protonated form interacts only to a small extent. Kinetic results suggest that the ligand is not involved in the rate determining step which is, in fact, associated with the release of a water molecule from the shell of the metal ions prior to complexation with the ligand. Enthalpy, entropy of activation, energy of the molecule and heat of formation were also calculated. A mechanism consistent with the kinetic data has been suggested.

**Keywords:** kinetics, Cu(II) ions,  $\gamma$ -aminobenzoic acid, rate constant, activation parameters.

### Introduction

$\gamma$ -aminobenzoic acid occurs in free state and in compounds other than folic acid in combination with a polypeptide of glutamic acid. It is also excreted in human urine in conjugated form with metal ions and in human faeces. It also cures hypoprothrombinaemia produced in young rats by feeding sulphasuxidine. It stimulates the growth of bacteria in the intestinal tract of humans thereby synthesise vitamin K necessary to restore the blood clotting mechanism. Therefore, the kinetic study of metal ions by  $\gamma$ -aminobenzoic would give important information regarding type of interaction of polypeptide of glutamic acid to metal ions. The structure of  $\gamma$ -aminobenzoic acid can be represented as:



( $\gamma$ -aminobenzoic acid) Zwitterion

According to Weber and Simeon eight membered rings are unstable due to large steric strain<sup>6</sup>. Thus it is obvious that reaction of Cu(II) with  $\gamma$ -aminobenzoic acid would proceed without *any* ring formation. The kinetics of complexation of metal ions Cu(II) by  $\gamma$ -aminobenzoic acid would give important information regarding the types of interaction of these ions with  $\gamma$ -aminobenzoic acid and the clearance of  $\gamma$ -aminobenzoic acid or metal ions in the form of metal-ligand complex through the animal body [1-5]. Such investigations are further desired for determining some important parameters, e.g. binding steps, rate constants corresponding to the binding steps  $K_{os}$  (outer sphere complex formation constant),  $k_o$  (Rate constant of water exchange) and activation parameters corresponding to the interaction of various reactive forms of the  $\gamma$ -aminobenzoic acid.

### Experimental method and materials

$\gamma$ -aminobenzoic acid (B.D.H.),  $\text{KNO}_3$  (B.D.H.) were used as such. Other chemicals used were of A.R. grade. The pH of ligand solution and metal ion solution was adjusted to same value

### Correspondence

**Yogeshwar Sharma**  
 Associate Professor, Department  
 of Chemistry, Motilal Nehru  
 College, Benito Juarez Marg,  
 University of Delhi South  
 Campus, New Delhi, India

using 2, 6-lutidine (Merck Schuchardt) and HCl. However, a slight change in pH value (~ 0.05 units) was observed after mixing of two solutions. The final pH was recorded from Radiometer pH meter, pH M26. pH's reported are those of reaction mixtures. The temperature of the system was maintained by immersion type thermostat (German NBE model). The kinetic runs were made on Aminco Morrow stopped flow spectrophotometer under pseudo first order conditions, i.e., [Cu(II)] >> [γ-aminobenzoic acid] at 620nm by pH indicator method. The total transmittance change was kept small using linear log photometer so that relative voltage change could be observed on the oscilloscope. Copper nitrate, 10<sup>-3</sup>M, buffer 10<sup>-2</sup>M Lutidine and, μ= 0.1 M KNO<sub>3</sub> was mixed with ligand solution and the traces from the oscilloscope gave excellent first order plots, from which second order rate constants (*k<sub>obs</sub>*) were computed by the relation

$$k_{obs} = k'_{obs}[Cu(II)] \tag{1}$$

Where *k'\_{obs}* is the pseudo first order rate constant.

**Results and Discussion**

**Complexation of Cu(II)- γ-aminobenzoic acid**

The kinetics of complexation of Cu(II)–γ-aminobenzoic acid was found to be of first order in copper ion, which was taken in large excess over ligand [8-9]. As protonated form of the ligand predominates in the pH range 2-5 ( pKa<sub>1</sub> = 2.50, pKa<sub>2</sub> = 4.87 ) [18], the kinetic study of interaction has been made in the pH range 2.01-3.14 at ionic strength 0.1M KNO<sub>3</sub> and at temperatures 25, 30, 35 and 40± 0.05 °C, under the condition [Cu(II)] >> [γ-aminobenzoic acid ]. Oscilloscope traces of voltage versus time were used to determine the values of pseudo-first order rate constants (*k'\_{obs}*), and these were further utilized to evaluate the values of second order rate constants (*k<sub>obs</sub>*), using equation (i). These rate constants are tabulated in Table 1.

**Table 1:** First order and Second order rate constants for the complexation of Cu(II) with γ-aminobenzoic acid at different pH's and temperatures

Temp. (±0.05°C)	pH	k'_{obs} x 10 <sup>-1</sup> (s <sup>-1</sup> )	k_{obs} x 10 <sup>-3</sup> (M <sup>-1</sup> s <sup>-1</sup> )	Temp. (±0.05°C)	pH	k'_{obs} x 10 <sup>-1</sup> (s <sup>-1</sup> )	k_{obs} x 10 <sup>-3</sup> (M <sup>-1</sup> s <sup>-1</sup> )
25	2.03	9.58	1.78	30	3.15	143	26.5
25	2.21	11.7	2.18	35	2.04	26.6	4.95
25	2.48	23.2	4.32	35	2.28	34.6	6.44
25	2.72	43.6	8.10	35	2.51	49.7	9.23
25	2.91	56.0	10.4	35	2.72	75.9	14.1
25	3.04	82.3	15.3	35	3.03	116	21.5
25	3.14	102	18.9	35	3.14	154	28.6
30	2.06	15.9	2.96	40	2.01	36.4	6.78
30	2.21	23.2	4.32	40	2.22	42.6	7.92
30	2.48	33.3	6.18	40	2.50	66.7	12.4
30	2.72	56.5	10.5	40	2.72	99.5	18.5
30	2.91	82.9	15.4	40	2.91	142	26.4
30	3.03	125	23.3	40	3.02	198	36.8


I = 0.10 M KNO<sub>3</sub>  
 [Cu(II)] = 5.38 x 10<sup>-2</sup> M  
 [γ-aminobenzoic acid ] = 4.86 x 10<sup>-3</sup> M

The rate equation for Cu(II) – γ-aminobenzoic acid interaction can be written as

$$\begin{aligned} \text{Rate} &= -d/dt[Cu(II)] \\ &= -d/dt[\gamma\text{-aminobenzoic acid}] \\ &= k_{obs} [\gamma\text{-aminobenzoic acid}] [Cu(II)] \tag{2} \\ &= k'_{obs} [\gamma\text{-aminobenzoic acid}] \tag{3} \end{aligned}$$

Where *k'\_{obs}*= *k<sub>obs</sub>* [Cu(II)]



The dissociation equilibria of γ-aminobenzoic acid can be represented as



$$K_1 = [H^+][HL] / [H_2L^+]$$

$$K_2 = [H^+][L^-] / [HL] \tag{5}$$

These three forms of ligand can react with Cu(II) ions as follows



Rate from equation 6, 7 and 8, can be written as

$$\begin{aligned} \text{Rate} &= d/dt[CuL^+] \\ &= \{ k_1 [H_2L^+] + k_2 [HL] + k_3 [L^-] \} [Cu(II)] \end{aligned} \tag{9}$$

From equation 5, it can be shown that

$$[L^-] = K_1 K_2 [H_2L^+] / [H^+]^2 \tag{10}$$

$$[HL] = K_1 [H_2L^+] / [H^+] \tag{11}$$

Substituting the values of [HL] and [L<sup>-</sup>] from equation (10) and (11) into equation (9) and on rearranging, we get:

$$\text{Rate} = \{ k_1 [H^+]^2 + k_1 K_1 [H^+] + k_3 K_1 K_2 \} [Cu(II)] [H_2L^+] / [H^+]^2 \tag{12}$$


Substituting the values of [HL] and [L<sup>-</sup>] from equation (10) and (11) into equation (2), and on simplification, we get:

$$\text{Rate} = k_{obs} \{ k_1 [H^+]^2 + K_1 [H^+] + K_1 K_2 \} [Cu(II)] [H_2L^+] / [H^+]^2 \tag{13}$$

Comparing equation (12) and (13) and assuming that diprotonated form [H<sub>2</sub>L<sup>+</sup>] is unreactive, i.e. k<sub>1</sub> ~ 0, it can be shown that

$$k_{obs} \{ [H^+]^2 + K_1 [H^+] + K_1 K_2 \} / K_1 [H^+] = k_2 + k_3 K_2 / K_1 [H^+] \tag{14}$$

Linear plots of  $k_{obs} \{ [H^+]^2 + K_1 [H^+] + K_1 K_2 \} / K_1 [H^+]$  versus  $[H^+]^{-1}$  at temperatures 25 30 35 and 40 ° C are shown in Fig.1. The values of K<sub>1</sub> and K<sub>2</sub> are obtained from the intercept and slope respectively. The values of K<sub>1</sub> and K<sub>2</sub> at 25°C were taken from literature and these values were corrected for different temperature using the equation (15)



**Fig 1:** Variation of  $k_{obs} \{ [H^+]^2 + K_1 [H^+] + K_1 K_2 \} \times 10^{-3} / K_1 [H^+]$  versus  $[H^+]^{-1}$  for Cu(II) -  $\gamma$ -aminobenzoic acid interaction at different temperatures


At the temperature of our investigation, the corrected values of  $K_3^T$  were evaluated using the thermodynamic relation:

$$pK_a^T = \{ \Delta H ( T_2 - T_1 ) / 4.576T_2 \cdot T_1 \} + pK_a^{25 \text{ } ^\circ\text{C}} \quad \text{-----(15)}$$

The values of  $k_2$  and  $k_3$  are reported in Table 2. Values of energies of activation corresponding to specific rate constants  $k_2$  and  $k_3$  were calculated from linear plot of  $\log k$  versus  $1/T$  and those of entropies and enthalpies of activation corresponding to  $k_2$  and  $k_3$  were calculated from linear plot of  $\log k/T$  versus  $1/T$

**Calculation of water exchange rate constant ( $k_o$ )**

The rate of exchange on metal ion is controlled by the rate of water exchange and rate determining step is the loss of water molecule from the inner coordination sphere, i.e.



The rate law in terms of outer sphere complex formation ( $K_{os}$ ) and rate constant of water loss ( $k_o$ ) from equation (19) be written as

$$\begin{aligned}
 d/dt (H_2O)_5ML^{(2-n)+} &= k_o [(H_2O)_5M^{2+}(H_2O)L^{n-}] \\
 &= k_o K_{os} [ M(H_2O)_6^{2+} ] [ L^{n-} ]
 \end{aligned} \quad \text{-----(17)}$$

So the rate law in terms of outer sphere complex formation ( $K_{os}$ ) for Cu(II)-  $\gamma$ -aminobenzoic acid can be written as

$$\text{Rate} = d /dt [Cu(II)L] = K_{os} k_o [Cu(II) ] [L] \quad \text{-----(18)}$$

Rate from equation 6, 7 and 8 can be written as

$$\begin{aligned}
 \text{Rate} &= d/dt[CuL^+] \\
 &= \{ k_1 [H_2L^+] + k_2 [HL] + k_3 [L^-] \} [Cu(II)]
 \end{aligned} \quad \text{-----(19)}$$

Assuming that the diprotonated form  $[H_2L^+]$  is unreactive  $k_1 \sim 0$ , and  $k_3 \gg k_2$  (Table 2), therefore equation (19) reduces to

$$\text{Rate} = k_3 [L^-][Cu(II)] \quad \text{-----(20)}$$

Comparing equation 18 and 20, we get

$$k_3 = K_{os} k_o \quad \text{-----(21)}$$

In order to calculate the value of  $k_o$ , we must know the value of  $K_{os}$ . The value of  $K_{os}$  was calculated using the equation which was given by Fuoss on statistical grounds,

$$K_{os} = 4\pi N a^3 \cdot e^{-u / K T} / 3000 \quad \text{----- (22)}$$

Where  $u = Z_1 Z_2 e_o^2 / aD - Z_1 Z_2 e_o^2 x / D ( 1 + xa)$

And  $x^2 = 8\pi N e_o^2 u / 1000DKT$


Where

- N = Avogadro's number
- a = Distance of closest approach of two ions
- K = Boltzmann's constant
- $e_o$  = Charge of an electron in esu units
- D = Bulk dielectric constant
- u = Ionic strength
- $Z_1 Z_2$  = Charge of reactants


Distance of closest approach of two ions often taken  $5\text{\AA}$  for reactions of two aqua cations with ordinary ligands. After substituting the values of all the terms in equation (22), the value of  $K_{os}$  can be approximated to  $1.98\text{ mol dm}^{-3}$  at all temperatures. The values of  $k_o$  are reported in Table 2. The high values of  $\Delta H^\ddagger$  corresponding to  $k_2$  step confirm that the mono-protonated form of  $\gamma$ -aminobenzoic acid is less reactive whereas low value of  $\Delta H^\ddagger$  corresponding to  $k_3$  supports the high reactivity of deprotonated form of  $\gamma$ -aminobenzoic acid. The negative value of entropy corresponding to  $k_3$  can be attributed to the fact that the transition state for this complex is highly charged and clearly shows that the reaction is between two oppositely charged ions (Table 3).

**Table 2:** Values of  $k_2$  and  $k_3$  and  $k_o$  for the complexation of Cu(II) with  $\gamma$ -aminobenzoic acid

Temp. ( $\pm 0.05^\circ\text{C}$ )	$k_2 \times 10^{-3}(\text{M}^{-1}\text{s}^{-1})$	$k_3 \times 10^{-6}(\text{M}^{-1}\text{s}^{-1})$	$k_o \times 10^{-6}(\text{s}^{-1})$
25	---	0.83	0.42
30	1.20	0.92	0.46
35	2.50	1.04	0.53
40	3.50	1.20	0.60



**Fig 2:** Variation of  $\log k_2$  and  $\log k_2 / T$  versus  $(1/T)$  for the interaction of Cu (II) - $\gamma$ -aminobenzoic acid




**Fig 3:** Variation of  $\log k_3$  and  $\log k_3 / T$  versus  $(1/T)$  on the interaction of Cu(II)- $\gamma$ -aminobenzoic acid

**Table 3:** Values of activation parameters corresponding to  $k_2$  and  $k_3$  steps for the complexation of Cu(II) with  $\gamma$ -aminobenzoic acid


	$k_2$	$k_3$
$\Delta H^\ddagger(\text{kJmol}^{-1})$	$74.1 \pm 2.9$	$14.3 \pm 1.1$
$\Delta E^\ddagger(\text{kJmol}^{-1})$	$81.0 \pm 1.7$	$15.9 \pm 0.6$
$\Delta S^\ddagger(\text{JK}^{-1}\text{mol}^{-1})$	$27.6 \pm 1.5$	$-108 \pm 4.5$

### Mechanism

The high values of  $\Delta H^\ddagger$  corresponding to  $k_2$  step confirm that the monoprotated form of  $\gamma$ -aminobenzoic acid is less reactive whereas low value of  $\Delta H^\ddagger$  corresponding to  $k_3$  supports the high reactivity of deprotonated form of  $\gamma$ -aminobenzoic acid. (Table 3). Due to strong electrostatic interaction between the positive charge of Cu(II) and negative charge on the oxygen, it is inferred that  $k_2$  is greater than  $k_3$ . The value of activation parameters corresponding to  $k_3$  further confirm that the deprotonated form is more reactive than the zwitterionic form of the ligand. This mechanism is further confirmed by the values of energy of activation and entropy of activation.



**Fig 4:** Deprotonated form of  $\gamma$ -aminobenzoic acid  
Final Geom Energy = -21526 kcal/mol  
Heat of Formation = 34745.2703 kcal/mol



**Fig 5:** Protonated form of  $\gamma$ -aminobenzoic acid  
Final Geom Energy = - 42524.18 kcal/mol  
Heat of Formation = - 26.7593 kcal/mol

This mechanism is further confirmed from the molecular modelling method in which the energy as well as the heat of formation of the zwitterionic form and protonated form has been calculated. Values of the energies as well as heat of formation were calculated after optimizing the geometry of the molecules<sup>[14-17]</sup>. These values are given in Fig 4 and 5. It has been found that the deprotonated form is more reactive than the protonated form.

#### References

1. Wilkins RG, Acc. Chem. Res. 1970; 3:408.
2. Siegel H, McCornik DB. J. Am. Chem. Soc., 1971; 43:2041.
3. Molinaro FS, Little RG, Jbers JA J. Am. Chem. Soc. 1977; 99:5628.
4. Eigen M, Wilkins RG. Adv. Chem. Ser. No. 1965; 49:55.
5. Eklund H *et al*, J. Mol. Bio. 1979; 1027:27.
6. Fast Reaction in Solution, E.F. Caldin, John Wiley and Sons, Inc., New York, 1964.
7. Voss RH, Jordan RB, J. Am. Chem. Soc. 1976; 98:2173.
8. Letter JE, Jordan RB. J. Am. Chem. Soc. 1975; 97:2381.
9. Malhotra HC, Jain AK, J. Ind. Chem. Soc. 1990; 67:365.
10. Malhotra HC, Sharma GC, Gazz. Chim. Ital. 1988; 118:323.
11. Malhotra HC, Yogeshwar Sharma, Jain AK, J. Ind. Chem. Soc. 1990; 67:192
12. Malhotra HC, Jai Prakash, Sharma GC, J. Ind. Chem. Soc. 1986; LX III:1045
13. Thompson MA, Zerner MC. J. Am. Chem. Soc., 1991; 113:8210.
14. Mark A, Thompson, Eric D, Glendening, David Feller J. Phys. Chem. 1994; 98:10465-10476.
15. Mark A Thompson, Gregory K Schenter J. Phys. Chem. 1995; 99:6374-6386.
16. Mark A Thompson, J. Phys. Chem. 1996; 100:14492-14507.
17. <http://www.arguslab.com/arguslab.com/ArgusLab.html>
18. <http://www.sas.upenn.edu/~marisa/documents/pkatables.pdf>