



ISSN (E): 2277-7695  
ISSN (P): 2349-8242  
NAAS Rating: 5.23  
TPI 2022; 11(11): 2231-2235  
© 2022 TPI  
[www.thepharmajournal.com](http://www.thepharmajournal.com)

Received: 04-08-2022  
Accepted: 09-09-2022

**Pawan Kumar Hooda**  
Department of Horticulture,  
School of Agriculture, Lovely  
Professional University,  
Phagwara, Punjab, India

**Jatinder Singh**  
Department of Horticulture,  
School of Agriculture, Lovely  
Professional University,  
Phagwara, Punjab, India

## Effect of edible coatings on jujube (*Zizyphus mauritiana* Lamk.) fruits: A review

**Pawan Kumar Hooda and Jatinder Singh**

### Abstract

Nowadays, Jujube fruits have tremendously marketed value because of its rich nutritive value. Jujube has a meager storage life (2-4 days) in ambient condition because of its highly perishable nature. The edible coating is an efficient approach to solve this problem as it contains edible substances like polysaccharide, proteins and lipids to protect the fruits. It is safe for the environment and human consumption. Nowadays, herbal edible coatings are utilized because of its nutritional value and benefit for human health. Edible coatings on jujube maintained the quality, increase the shelf life and prevent microbial growth in both ambient and cold storage condition.

**Keywords:** Ber, jujube, edible coating, quality, shelf life

### Introduction

Jujube (*Zizyphus mauritiana* Lamk.) is the most valuable fruit crop in tropical and sub-tropical regions (Baloda *et al.* 2012) [6]. The native place of *Z. mauritiana* is central Asia. It belongs to the family Rhamnaceae. Ber fruits are a rich source of vitamins as well as carotenoids, pectin and mineral elements. It is noteworthy that ber fruits have an ascorbic acid (vitamin C) content higher than citrus and second only to guava among tropical fruits. Indian jujubes are cultivated in several Asian countries, mainly China, Pakistan, Thailand, Korea, Iran, and Vietnam. India is the world's top producer and this species is mainly present in its arid and semi-arid regions. The pulp of ber with high nutrient value has been comprehensively perceived and it contains varied phytochemicals like flavonoids, sugars, phenolic acids and nutrients like vitamin A and C, P, Ca, and Fe. It is considered to be the poor man's apple which contains good quantities of vitamins, minerals and sugar. Ber has been generally viewed as a useful organic product in India, unique parts and delightful taste. In India area is roughly one lakh ha and the yearly production of jujube is evaluated 10 lakh ton and productivity of jujube is 10 metric ton ha<sup>-1</sup> and India positions 2nd among jujube-developing nations after China. The significant territories for jujube development in India are Gujarat, Uttar Pradesh, Maharashtra, Tamil Nadu, West Bengal, Andhra Pradesh, Jammu and Kashmir, Punjab, Madhya Pradesh, Karnataka, Rajasthan, and Haryana (Bal *et al.* 2014) [4]. The growing areas are given in the following:-

### State Growing Areas

Punjab – Patiala, Sangrur, Muktsar  
Rajasthan – Jaipur, Jodhpur, Bharatpur  
Haryana – Hisar, Rohtak, Jind, Panipat, Mahendergarh  
Karnataka – Gulbarga, Belagaum, Raichur Bijapur, Bellary  
Gujarat – Banaskantha, Sabarmati  
Tamil Nadu – Salem, Ramanathapuram, Tirunelveli, Dharmapuri

Ber is perishable and has a storage life (2-4 days) at room storage (Meen *et al.* 2009) [24]. Fresh jujube increase physiological weight loss and rot because of high respiration rate and other enzymatic activity during storage (Wang *et al.* 2011) [41] and is a non-climacteric fruit and maturity is judged by the appearance of fruit peel color (Saranant *et al.* 1992) [38]. The color changes of the ber from bright green to red and black due to quinone polymerization and anthocyanin oxidization under improper storage conditions. Several strategies which include refrigeration changed surroundings storage, chemical preservatives and packaging are getting used to limit deleterious outcomes (Zhang and Quantick *et al.* 1997) [34].

**Corresponding Author:**  
**Jatinder Singh**  
Department of Horticulture,  
School of Agriculture, Lovely  
Professional University,  
Phagwara, Punjab, India

For instance, expert long covering postponed the maturity of kalpataru fruit by changing the inner environments (Banks *et al.* 1984) [7], and comparable outcomes were gotten in Malus and Pyrus fruits.

(Quantick & Zhang *et al.* 2007) [33] found that a consumable covering dependent on sucrose esters of unsaturated fats deferred pericarp caramelizing of litchi organic product.

(Lachman *et al.* 1986) [20] and (Lowe *et al.* 1963) [21] thin layer of edible materials restricted the loss of water and respiration rate of food. Herbs like aloe vera, cinnamon oil, ginger oil are used as edible coatings because of their nutritional (vitamins,

minerals) and medicinal values (Chauhan *et al.* 2014) [9], (Martinez *et al.* 2006) [23], (Nasution *et al.* 2015) [28] and herbs also contain some antimicrobial activity (Douglas *et al.* 2005) [14]. It is safely eaten with the food product (Anand *et al.* 2007) and coatings have long been regarded to guard perishable food merchandise against deterioration.

The edible coating contains numerous materials like gums, proteins, lipids, or their mixtures, which extending the storage life of post-harvested vegetables and fruits (Raghav *et al.* 2016) [35].




Fig: Various types of edible coating

### 1. Efficacy of edible coatings on quality and shelf life of Jujube

The efficacy of edible coatings in maximize the storage-periods of fresh fruits by retaining quality. The efficacy of edible coatings (1-MCP & chitosan) on the quality and enzyme activities of jujube fruit for stored 42 days at  $0 \pm 1$  °C. The result revealed that individual treatment of 1-MCP

and chitosan reduced the rotting rate, firmness, soluble solids and increased the enzyme activities of peroxidase, superoxide dismutase, ascorbate peroxidase and catalase. The combined treatment of 1-MCP and chitosan was more efficacy for postharvest quality and storage life than individual treatments (Cheng *et al.* 2020) [11]. Both carnauba wax and carnauba wax + glycerol monolaurate coatings were reduced the water loss,

ethylene production and respiration rate, maintain pectin methylesterase, cellulose, polygalacturonase activities and also maintained the skin colour and chlorophyll content and vitamin C in comparison to the control. The combined coating of CW+Gradient multilayer (GML) had increased shelf life and maintained quality when stored for 12 days at 20°C (Chen *et al.* 2019) [10]. Evaluation of the effect of surface coatings and packaging materials on ascorbic acid and sugars of apple ber (*Zizyphus mauritiana*) at different storage conditions. The result showed coating of Chitosan 1% with High density polyethylene (HDP) plus cold storage (10±2°C) extent the shelf life upto 21 days and also increased vitamin C and total sugar than the all treatments (Bhavana *et al.* 2019) [8].

The efficacy of edible coatings CaCl<sub>2</sub> + Palm oil methylester (PME) on storage quality of jujube fruit. The results revealed that CaCl<sub>2</sub> (1% w/w) combined with pectin methylesterase (15U/mL) under vacuum impregnation, delayed physiological weight loss and increasing quality of fruits by retaining Firmness, vitamin C and soluble solids content (Zhang *et al.* 2019) [46]. The influence of edible coatings i.e. aloe vera gel (33 and 50% v/v), carboxymethyl cellulose and pectin (1, 1.5 and 2% w/v) on postharvest quality of fresh Chinese jujube fruits during refrigerated storage at 4°C for 40 days. The results showed that both aloe vera gel 30% and pectin 1.5% coated fruits decreased physiological weight loss upto 30%, when stored for 40 days but higher sensory quality and TA observed fruits coated with 50% v/v Aloe vera, 1 and 2% w/v carboxymethyl cellulose than control condition. (Moradinezhad *et al.* 2018) [26] observed both vera gel (33 and 50% v/v), carboxymethyl cellulose and pectin (1, 1.5% w/v) were maintained quality of ber fruits when storage at 4°C for 40 days.

The combined mixture of guar gum with aloe-vera minimize the loss of physiological weight, titratable acidity, decay, firmness of fruit and also maintain ascorbic acid & greenish-yellow colour of the fruit upto 15 day at ambient storage condition (Arghya *et al.* 2018) [1]. The study evaluated by dipping of ber in hydros solution of C<sub>2</sub>H<sub>2</sub>O<sub>4</sub> for 10 minutes at varied concentrations (2 mM, 4 mM, 6 mM, 8 mM and 10 mM). (Ravi *et al.* 2018) [37] revealed that 10mM concentration was more effective by maintaining physiology & enzymatic activity and also extended the storage period up to 9 days in ambient condition other than all concentrations. Five treatments (T<sub>1</sub>- GA3 40 ppm, T<sub>2</sub> - CaCl<sub>2</sub> 1.0%, T<sub>3</sub>- Aloe vera Gel, T<sub>4</sub> - Olive oil, T<sub>5</sub> - Almond oil) were coated on ber and store at two different conditions (ambient and cold storage). The results showed CaCl<sub>2</sub> 1.0% coating was more effective in cold storage (shelf life 30 days) and 9 days at ambient condition but maximum total soluble solid was observed in GA<sub>3</sub> coated @ 40 ppm in cold storage Jain *et al.* (2017) [18].

The effects of three treatments T<sub>1</sub> = control, T<sub>2</sub>= corn starch (1%) and T<sub>3</sub>= aloe vera gel (2%) on jujube. The result indicated aloe-vera coating was effective to maintain quality by reducing the ripening rate, minimize the loss of ascorbic acid and also reduce shrinkage percentage, lower physiological loss (weight) and maximum colour retention in comparison to uncoated ones. Corn with aloe vera gel coated fruits had good shelf life (Mani *et al.* 2017) [22]. The effects of edible coatings (1% pullulan, 1% chitosan and 2% CaCl<sub>2</sub>) on quality and enzyme activity of jujube fruit during storage. The results showed all three coatings (1% pullulan, 1% chitosan and 2% CaCl<sub>2</sub>) retained the fruit quality and antioxidant activity at storage condition. The observation was that among

these three treatments 2% CaCl<sub>2</sub> was best, followed by other two coatings (Kou *et al.* 2017) [19]. Using coating with varied concentrations of tea polyphenols (1, 3, and 5 g/L) integrated with alginate-based edible coating for retain quality of jujube under room temperature for 8 days. The alginate-based edible coating mixed with Tea polyphenols (1g/L) decreased respiration rate, red indices, chlorophylls content, malonaldehyde content and also maintained the activities of antioxidant enzymes, vitamin C, and total phenol content under ambient storage condition (Zhang *et al.* 2016) [45].

The efficiency of coatings of alginate (2%)-olive oil (0.2%) combined with antioxidants on the quality and storage period of jujube at ambient condition with 65% relative humidity. The combined treatments decreased the physiological weight loss, increase TSS and also maintained quantity of antioxidants by enhancing the storage life of the ber fruit (Rao *et al.* 2016) [36]. Increasing the shelf-life and quality jujube fruits (BAU variety) by using different edible coatings like gum tragacanth (1.0%, 1.5%, 2.0%), guar gum (1%, 1.5%, 2%), chitosan (0.5%, 1.0%, 2.0%). The edible coatings decrease respiration rate and delayed the decrease of TSS, vitamin C, titratable acidity, total and reducing sugar, weight loss and maintain quality & green colour in jujube fruit as compared to uncoated fruit. Guar gum (2%) extended the storage life of jujube up to 16 days (Dutta *et al.* 2016) [15]. 1% chitosan combined with 0.1% cinnamon oil, maintain quality by reducing the weight loss and decay of fruits when stored at 4°C for 60 days but reduction occurred in vitamin C and titratable acid (Xing *et al.* 2015) [42].

The combination of chitosan coating with UV radiation on maintenance of jujube under room temperature. The results revealed delayed increase of the respiration rate, weight loss, firmness, and maintain decrease of vitamin C, chlorophyll content, and activity of (superoxide dismutase, catalase, and Peroxidase). This treatment of ultraviolet irradiation with chitosan was best to retain the postharvest quality of ber fruit (Zhang *et al.* 2014) [47]. The effects of chitosan 1% with nano-silicon dioxide 0.04% on the storage quality for 32 days at ambient condition. The results revealed that this composite coating reduced the physiological weight loss, spoilage, respiration rate, red index and also increase the enzyme activities i.e. catalase, superoxide dismutase, and peroxidase (Yu *et al.* 2012) [44]. The efficacy of chitosan and 1-MCP coating on quality evaluation of jujube fruit. The results indicated combine of 1-MCP (600 nl l<sup>-1</sup>) with chitosan (1.5 g/100 ml) reduced respiration rate, climacteric ethylene evolution stem-end rots incidence, & decreased PG and LOX activities and also maintained chlorophyll content, vitamin c, firmness and TSS of ber fruits. This combined treatment 1-MCP and chitosan extend shelf life and maintain quality of jujube fruit (Yu *et al.* 2012) [44].

## 2. Efficacy of edible coatings against Spoilage Microorganisms

Edible coatings in extending the storage-life of fresh fruits by inhibiting the growth of spoilage bacteria. The combinational efficacy of essential oils to control decaying caused fungus i.e. (*Alternaria alternata* and *Penicillium expansum*) and to maintain fruit quality. The results showed combinational treatment of cinnamon (0.156g L<sup>-1</sup>), thyme (0.078 g L<sup>-1</sup>) and rosemary (0.625g L<sup>-1</sup>) restrict both spoilage fungus i.e. *Alternaria alternata* and *Penicillium expansum* in ber fruit. The essential oils treated fruits were reduce physiological



weight loss, respiration rate and also retained vitamin-C (Nikkhah, & Hashemi *et al.* 2020) [29]. The effects of coatings based chitosan (0.1 mol L<sup>-1</sup>) incorporated with sodium silicate (hundred mmol L<sup>-1</sup>) to induces resistance against spoil caused by fungus during postharvest storage. The results indicated that combined treatment of chitosan with sodium silicate, decreased the physiological weight loss, lesion diameter, decay, red index and increased some substances like vitamin C, flavonoids, lignin content and total phenolic compounds. Chitosan + sodium silicate induced the activities of superoxide dismutase, phenylalanine ammonia lyase, polyphenol oxidase, β-1, 3-glucanase and peroxidase, chitinase which accumulate the disease resistance of jujube (Guo *et al.* 2019) [16]. The edible coatings along with potassium meta bisulphate reduced the microbial activity (both bacteria and fungi) (Padmaja and John *et al.* 2014) [30]. Chitosan application inhibited germination of spore, germ tube length and mycelial growth by causing plasma membrane collapse of *P. expansum* and also restricting postharvest diseases without any harmful effect on fruit quality (Wang *et al.* 2014) [44, 47]. Chitosan mixed with cinnamon oil 2.0%, completely reduced the growth activity of blue mold fungus (*Penicillium citrinum*) and also retained ascorbic acid and phenolic compounds in jujube (Xing *et al.* 2011) [43].

### Conclusion

Palatable coatings are utilized by food industry from many years to store vegetables and fruits. Varied types of edible coatings are used like polysaccharide, proteins and lipids. Nowadays herbal edible coatings because of its environment friendly and safety for consume with fruits. This review indicates edible coatings increases the shelf life (by reducing physiological weight loss, respiration rate) and maintains quality and also prevent microbial growth of ber fruit at storage condition.

### References

1. Arghya M, Prasanna VSSV, Shuvadeep H, Praveena J. Efficacy of edible coatings blended with aloe vera in retaining post-harvest quality and improving storage attributes in Ber (*Zizyphus mauritiana* Lamk.). International journal of chemical studies. 2018;6(6):1727-1733.
2. Ayranci. The effect of edible coatings on water and vitamin C loss of apricots (*Armeniaca vulgaris* Lam.) and green peppers (*Capsicum annuum* L.). Food Chemistry. 2004;87(3):339-342.
3. Azarakhsh N, Osman A, Ghazali HM, Tan CP, Mohd Adzahan N. Lemongrass essential oil incorporated into alginate-based edible coating for shelf-life extension and quality retention of fresh-cut pineapple. Postharvest Biology and Technology. 2014;88:1-7.
4. Bal JS. Development and production of Indian jujube (Ber) in India. In XXIX International Horticultural Congress on Horticulture: Sustaining Lives, Livelihoods and Landscapes (IHC2014). 2014;3(1116):15-22.
5. Baldwin EA, Burns JK, Kazokas W, Brecht JK, Hagenmaier RD, Bender RJ, *et al.* Effect of two edible coatings with different permeability characteristics on mango (*Mangifera indica* L.) ripening during storage. Postharvest Postharvest Biology and Technology. 1999;17:215-226.
6. Baloda S, Sehrawat SK, Yadav BS, Ahlawat VP, Singh S. Present status of ber production and future thrusts in India-a review. Agricultural Reviews. 2012;33(3):256-264.
7. Banks NH. Some effects of TAL Pro-long coating on ripening bananas. Journal of Experimental Botany. 1984;35(1):127-137.
8. Bhavana VJ, Sreedhar M, Mishra RP. Effect of surface coatings and packaging materials on ascorbic acid and sugars of apple ber (*Zizyphus mauritiana*) at different storage conditions. Journal of entomology and zoology studies. 2019;7(2):277-282.
9. Chauhan S, Gupta KC, Agrawal M. Application of Biodegradable Aloe vera gel to control post-harvest decay and longer the shelf life of Grapes. International Journal Current Microbiology and Applied Sciences. 2014;3(3):632-642.
10. Chen H, Sun Z, Yang H. Effect of carnauba wax-based coating containing glycerol monolaurate on the quality maintenance and shelf-life of Indian jujube (*Zizyphus mauritiana* Lamk.) fruit during storage. Scientia horticulturae. 2019;244:157-164.
11. Cheng S, Yu Y, Guo J, Chen G, Guo M. Effect of 1-methylcyclopropene and chitosan treatment on the storage quality of jujube fruit and its related enzyme activities. Scientia Horticulture. 2020;265:109281.
12. Choi SH, Ahn JB, Kozukue N, Levin CE, Friedman M. Distribution of free amino acids, flavonoids, total phenolics, and antioxidative activities of jujube (*Zizyphus jujuba*) fruits and seeds harvested from plants grown in Korean. Journal of Agriculture Food Chemistry. 2011;59:6594-6604.
13. Davies HD, Elson CU, and M Hayes ER. N10-carboxymethyl Chitosan, a New Water Soluble Chitosan Derivative. 4th International Conf. on Chitosan and Chitosan. Trondheim, Norway; c1988.
14. Douglas JH, Smallfield B. Herb spice and essential oil: post-harvest operation in developing country, 2<sup>nd</sup> Edition; c2005. p. 45-55.
15. Dutta P, Dey K, Ghosh A, Bhowmick N, Ghosh A. Effect of edible coatings for enhancing shelf- life and quality in Ber (*Zizyphus mauritiana* Lamk.) fruits. Journal of Applied and Natural Science. 2016;8(3):1421-1426.
16. Guo Y, Zhou J, Zhang J, Zhang S. Chitosan combined with sodium silicate treatment induces resistance against rot caused by *Alternaria alternata* in postharvest jujube fruit. Journal of Phytopathology. 2019;167(7-8):451-460.
17. Holton TA, Cornish EC. Genetics and biochemistry of anthocyanin biosynthesis. The Plant Cell. 1995;7(7):1071.
18. Jain D, Kachwaya DS, Kuchi VS, Vikas G. Influence of post-harvest treatments on storage behaviour and fruit value of ber (*Zizyphus mauritiana*) cv. gola. Plant Archives. 2017;17(2):1277-1282.
19. Kou X, Li Y, Wu J, Chen Q, Xue Z. Effects of edible coatings on quality and antioxidant activity of *Zizyphus Jujuba* Miller cv. Dongzao during storage. Transactions of Tianjin University. 2017;23(1):51-61.
20. Lachman L, Lieberman HA, Kanig JL. The Theory and Practice of Industrial Pharmacy. Lea and Febiger Publishing Co.; Philadelphia; c1986. p. 348-373.
21. Lowe E, Durkee EL, Hamilton WE, Watters GG, Morgan AI. Continuous raisin coater. Food Technol.

- 1963;11(2):109-111.
22. Mani A, Jain N, Singh AK, Sinha M. Effects of aloevera edible coating on quality and postharvest physiology of ber (*Zizyphus Mauritiana* Lamk.) under ambient storage conditions. *Annals of Horticulture*. 2017;10(2):138-146.
  23. Martínez-Romero D, Alburquerque N, Valverde JM, Guillén F, Castillo S, Valero D, *et al.* Postharvest sweet cherry quality and safety maintenance by Aloe vera treatment: a new edible coating. *Postharvest Biology and Technology*. 2006;39(1):93-100.
  24. Meena HR, Kingsly ARP, Jain RK. Effect of post-harvest treatments on shelf life of ber fruits. *Indian Journal of Horticulture*. 2009;66(1):58-61.
  25. Shin, Melvin. The application of edible polymeric films and coatings in the food industry. *Journal of Food Processing and Technology*. 2012;4(2):1-2.
  26. Moradinezhad F, Naeimi A, Farhangfar H. Influence of edible coatings on postharvest quality of fresh Chinese jujube fruits during refrigerated storage. *Journal of Horticulture and Postharvest Research*. 2018;1(1):1-14.
  27. Morton J. Indian Jujube. In: *Fruits of Warm Climates*. (Ed. Morton, J.F.) Miami, Florida; c1987. p. 272-275. <http://www.hort.purdue.edu/newcrop/morton/Indianjujub e>.
  28. Nasution Z, Ye JNW, Hamzah Y. Characteristics of fresh-cut guava coated with aloe vera gel as affected by different additives. *Agriculture and Natural Resources*. 2015;49(1):111-121.
  29. Nikkhah M, Hashemi M. Boosting antifungal effect of essential oils using combination approach as an efficient strategy to control postharvest spoilage and preserving the jujube fruit quality. *Postharvest Biology and Technology*. 2015;164:111159.
  30. Padmaja N, Bosco SJD. Preservation of jujube fruits by edible Aloe vera gel coating to maintain quality and safety. *Indian Journal of Scientific Research and Technology*. 2014;2(3):79-88.
  31. Pareek OP. Indian Jujube and Pomegranate. In: *50 Years of Crop Science Research in India*. (Eds. Paroda RS, Chadha KL). Indian Council of Agricultural Research, New Delhi; c1997. p. 557-565.
  32. Qiuping Z, Wenshui X. Effect of 1- methylcyclopropene and/or chitosan coating treatments on storage life and quality maintenance of Indian jujube fruit. *LWT-Food Science and Technology*. 2007;40(3):404-411.
  33. Quantick, Zhang. Effects of chitosan coating on enzymatic browning and decay during postharvest storage of litchi (*Litchi chinensis* Sonn.) fruit. *Postharvest Biology and Technology*. 1997;12(2):195-202.
  34. Raghav PK, Agarwal N, Saini M. Edible coating of fruits and vegetables: A review. *International journal of scientific research and modern education*. 2016;1(1):188-204.
  35. Rao TR, Baraiya NS, Vyas PB, Patel DM. Composite coating of alginate-olive oil enriched with antioxidants enhances postharvest quality and shelf life of Ber fruit (*Zizyphus mauritiana* Lamk. Var. Gola). *Journal of food science and technology*. 2016;53(1):748-756.
  36. Ravi K, Pareek S, Kaushik RA, Ameta KD. Effect of oxalic acid on ripening attributes of 'Gola'ber (*Zizyphus mauritiana* L amk.) fruit during storage. *International Journal of Chemical Studies*. 2018;6(5):403-408.
  37. Saranant S. Status of the tropical fruit industry in Thailand *Acta Horticulture*. 1992;292:13-23.
  38. Voilley, Debeaufort FJ, Quezada-Gallo A. Edible films and coatings: tomorrow's packaging: a review. *Crit Rev Food Sci Nutr*. 1998;38:299-313.
  39. Wang L, Wu H, Qin G, Meng X. Chitosan disrupts *Penicillium expansum* and controls postharvest blue mold of jujube fruit. *Food Control*. 2014;41:56-62.
  40. Wang YF, Tang F, Xia JD, Yu T, Wang J, Azhati R, Zheng XD. A combination of marine yeast and food additive enhances preventive effects on postharvest decay of jujubes (*Zizyphus jujuba*). *Food Chemistry*. 2011;125:835-840.
  41. Xing Y, Lin H, Cao D, Xu Q, Han W, Wang R, *et al.* Effect of chitosan coating with cinnamon oil on the quality and physiological attributes of China jujube fruits. *BioMed research international*; c2015.
  42. Xing Y, Xu Q, Che Z, Li, X, Li W. Effects of chitosan-oil coating on blue mold disease and quality attributes of jujube fruits. *Food & Function*. 2011;2(8):466-474.
  43. Yu Y, Zhang S, Ren Y, Li H, Zhang X, Di J. Jujube preservation using chitosan film with nano-silicon dioxide. *Journal of Food Engineering*. 2012;113(3):408-414.
  44. Zhang L, Li S, Dong Y, Zhi H, Zong W. Tea polyphenols incorporated into alginate-based edible coating for quality maintenance of Chinese winter jujube under ambient temperature. *LWT-Food Science and Technology*. 2016;70:155-161.
  45. Zhang L, Wang P, Chen F, Lai S, Yu H, Yang H. Effects of calcium and pectin methylesterase on quality attributes and pectin morphology of jujube fruit under vacuum impregnation during storage. *Food chemistry*. 2019;289:40-48.
  46. Zhang S, Yu Y, Xiao C, Wang X, Lei Y. Effect of ultraviolet irradiation combined with chitosan coating on preservation of jujube under ambient temperature. *LWT-Food Science and Technology*. 2014;57(2):749-754.