



ISSN (E): 2277-7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2022; 11(11): 498-501
© 2022 TPI
www.thepharmajournal.com
Received: 30-08-2022
Accepted: 06-10-2022

M Lakshmi Naga Nandini
Ph.D. Scholar, Department of
Plant Pathology, College of
Horticulture, Dr. YSR
Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

B Srinivasulu
Director of Extension, Dr. YSR
Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

K Gopal
Registrar, Dr. YSR
Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

Dr. Ch Ruth
Professor, Dr. YSR
Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

P Rama Devi
³Professor, COH, Dr. YSR
Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

M Ravindra Babu
Senior Scientist, Horticultural
Research Station, Dr. YSR
Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

VV Padmaja
Assistant Professor, COH, Dr.
YSR Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

Corresponding Author:
M Lakshmi Naga Nandini
Ph.D. Scholar, Department of
Plant Pathology, College of
Horticulture, Dr. YSR
Horticultural University,
Venkataramannagudem,
Andhra Pradesh, India

Effect of date of planting on pest and viral disease incidence on bell pepper (*Capsicum annuum* var *grossum* Sendt.) under naturally ventilated polyhouse

M Lakshmi Naga Nandini, Dr. B Srinivasulu, Dr. K Gopal, Dr. Ch Ruth, P Rama Devi, Dr. M Ravindra Babu and Dr. VV Padmaja

Abstract

The effect of three dates of planting viz. 1st December, 1st January and 1st February on pest and viral diseases incidence of bell pepper variety Arka Mohini was studied in an experimental field at West Godavari district in Andhra Pradesh for two consecutive seasons (2020-21 & 2021-22). The five insect pests namely thrips (*Scirtothrips dorsalis* Hood), aphids (*Aphis gossypii* Glover. and *Myzus persicae* Sulzer), whitefly (*Bemisia tabaci* Gennadius), yellow mite (*Polyphagotarsonemus latus* Banks) and fruit borer (*Helicoverpa armigera* Hubner) and viral diseases like chilli leaf curl and capsicum chlorosis were recorded.

Crop planted on 1st December showed highest average population of thrips (1.24/5 leaves), aphids (2.09/5 shoots), whitefly (2.57/5 leaves), fruit borers (0.97/plant), capsicum chlorosis virus incidence (18.67%) and chilli leaf curl virus incidence (34.58%) but lowest average population of yellow mite (0.42/5 leaves) and damage of plant due to mite (15.59%). Crop planted on 1st January exhibited moderate population of thrips (0.76/5 leaves), aphids (1.44/5 shoots), whitefly (2.33/5 leaves) and fruit borers (0.73/plant), capsicum chlorosis virus incidence (10.92%) and chilli leaf curl virus incidence (26.70%) except yellow mite (0.71/5 leaves). Pest and virus incidence was lowest on 1st February planted crop. Crop planted on 1st February gave highest yield of marketable fruits (18.3t/ha) followed by 1st January (15.2 t/ha) and 1st December (13.3t/ha) planted crop.

Keywords: Bell pepper, planting dates, pest incidence, leaf curl virus, capsicum chlorosis virus

Introduction

Capsicum (*Capsicum annuum* var *grossum* Sendt.), also known as 'Bell pepper' or 'Sweet pepper', is a highly remunerative vegetable crop grown in most parts of the world. India contributes one fourth of world production of capsicum with an annual production of 0.9 million tons from about 0.85-million-hectare area [1]. In India, Karnataka has highest area under capsicum cultivation (3.10 thousand hectares) with a production of 45.80 thousand metric tons [2]. In Andhra Pradesh the crop is grown over small areas. Owing to its increasing demand especially by urban consumers, the farmers are showing increased interest in growing the crop in open fields and under protected structures mainly during winter months, and both area and production has increased considerably in recent years. Moreover, it has high potential for export. About 35 species of insect and mite pests have been reported to infest bell pepper, of which thrips (*Scirtothrips dorsalis* Hood and *Thrips palmi* army), aphids (*Aphis gossypii* Glover and *Myzus persicae* Sulzer), whitefly (*Bemisia tabaci* Gennadius), capsule borers (*Helicoverpa armigera* Hubner and *Spodoptera litura* Fabr.) and yellow mite (*Polyphagotarsonemus latus* Banks) cause serious damage to the crop in different regions of India [3,4]. Ahmed (2005) [5] has estimated the crop losses up to 34 per cent due to attack of sucking pests. In addition to de-sapping, sucking pests also transmit viral diseases, which become a limiting factor in the cultivation of capsicum [6]. During the past decade, white fly has become a prominent problem worldwide especially in the subtropical agro ecosystems. Millions of dollars have been lost as a result of direct damage, honeydew contamination, and fungal growth and virus diseases caused by white fly -transmitted geminiviruses [7,8].

As pesticide residues in bell pepper are of great concern from the point of view of exports and domestic consumption as well, nonchemical pest management strategy like adjusting the date of planting for avoiding the pests and viruses without sacrificing the yield may be a better approach. Hence, an attempt was made to study the impact of different dates of planting of capsicum on pest and capsicum chlorosis virus and chilli leaf curl virus incidence keeping in mind the cropping sequence of the concerned area.

Materials and Method

A Field experiment of capsicum varieties Arka Mohini and Arka Atulya was conducted in an experimental field at West Godavari district in Andhra Pradesh during Rabi 2020-21 and 2021-22. The area falls in the Krishna-Godavari zone of Andhra Pradesh and lies between 16°63' N latitude and 81°27' E longitude with mean annual rainfall of 900 mm, maximum temperature around 41^o C during May and minimum temperature around 18^o C during January. The relative humidity of the area lies between 50% in March and 90% in July.

The capsicum variety Arka Mohini was grown in 24 plots (6 treatments and 4 replications) each measuring 5.4 sq m and following the design RBD. Thirty days old seeding was planted at spacing of 60cm X 45 cm. on three different dates, i.e., 1st December, 1st January and 1st February each year. Tenthon organic manure, N: P₂O₅: K₂O @ 100:80:80 kg /ha and micronutrient mixture of copper, zinc, boron @ 15kg /ha applied into soil at the time of transplanting. After that 10 kg/ha molybdenum and sulphur mixture was applied in soil at the time of earthing up. Apart from this capsicum crop was sprayed with water soluble fertilizer like N: P₂O₅:K₂O: 19:19:19 at every three weeks interval after 2 months of transplanting @ 3g/lit of water. Weeding up was done before earthing up. Then earthing up was completed after 20 days of

transplanting followed by first irrigation. After that the crop was irrigated 8 to 10 days interval throughout cropping period. The crop was raised / grown as per the recommended package of practices except pesticide measure.

All the observations were recorded during morning hours between 6:30 AM to 8:30 AM with the help of a hand lens (10X). The pest incidence was recorded from five fixed plants/plot, selected randomly for this purpose, at 10 days interval starting from 15 days after planting. The population of whiteflies (nymphs and adults) was recorded from five leaves one each from the upper, middle and lower position from each plant. Thrips population was recorded from five twigs of each plant by shaking over a piece of wet white coloured cloth. The population of aphids was recorded from five numbers of 10 cm terminal shoot from each plant. Mite count was recorded from five terminal leaves on each plant. Population of lepidopteron larvae of the fruit borer was recorded on the whole plant basis. More over the damage index by chilli leaf curl virus was recorded in 1-5 scale following (Table-1) [9]. The percentage of fruits damaged by fruit borer, and fruit yield were recorded at harvest. Data obtained were subjected to analysis of variance after suitable transformation. In case of percentage observations pooled data analysis was obtained after angular transformation.

Table 1: Scoring procedure for leaf curl virus damage

Score	Symptoms
0.	No symptoms
1	1 to 25% leaves per plant showing curling or damage
2	26 to 50% leaves per plant showing curling moderately damage
3	51 to 75% leaves per plant showing curling, heavily damaged, malformation of growing points, and reduction in plant height.
4	>75% leaves per plant showing curling, severe and complete destruction of growing points, drastic reduction in plant height, defoliation and severe malformation

Result and Discussion

During the course of study, five pest species viz. thrips, aphid, whitefly, fruit borer and yellow mite were detected causing damage to bell pepper at various growth stages of the crop. Table 2 represented that, during the first date planting (1st December), the thrips population was ranged from 1.81 to 1.96 per 5 leaves with a calculated pooled value of 1.24 per 5 leaves. The aphid population was varied from 3.46 to 5.45 per 5 shoots with calculated pooled population of 2.09 per 5 shoots. The white fly population was ranged from 5.67 to 7.65 per 5 leaves with calculated pooled value of 2.57 per 5 leaves. The yellow mite population was recorded from 0.15 to 0.22 per 5 leaves with calculated pooled value 0.42 per 5 leaves. Finally, fruit borer population was ranged from 0.91 to 1.01 per plant with pooled data 0.97 per plant though out the crop growth period. The similar result was found by Pathipati, V.L. (2015) [10] who reported that the population and damage levels, per cent yield loss of bell pepper due to insect pests and insecticide usage pattern were recorded in open field and poly house conditions in and around Hyderabad. The insect pests, viz., thrips, mite, cut worm, blossom midge, and fruit borer, *S. litura* incidence were recorded, whereas, in poly house in addition to the above pests, aphids and whiteflies incidence were also recorded. The present survey is in line with the findings of Sunitha (2007) [3] who took a fixed plot survey to record the natural incidence of insect pests of capsicum in major growing areas in and around Dharwad and Belgaum, Karnataka. The survey revealed the occurrence of

cutworm, *Agrotis ipsilon* (Hufn.) (30 to 40%), aphids, *M. persicae* (0.00 to 1.6 nymphs and adults/ 3 leaves), thrips, *S. dorsalis* (0.00 to 2.00 nymphs and adults/ leaves) and fruit borer, *Helicoverpa armigera* (Hub) (20.68 to 26.16% damage), respectively.

In second date planting (1st January), it was observed that thrips population ranged from 0.67 to 0.51 per 5 leaves with pooled data 0.76 per 5 leaves, aphid population varied from 1.36 to 2.96 per 5 shoots with pooled population 1.44 per 5 shoots, white fly from 4.70 to 6.23 per 5 leaves with pooled data 2.33 per 5 leaves, yellow mites from 0.45 to 0.57 per 5 leaves with pooled data 0.71 per 5 leaves and fruit borer ranged from 0.44 to 0.63 per plant with pooled data 0.73 per plant though out the growth period. Similar results were also obtained by Meena *et.al.* (2013) [11] who observed that the incidence of thrips, whiteflies, aphids and mites were appeared on the chilli crop soon after transplanting, while the aphid appeared little late during both the years.

In case of 1st February plated crop, it was noted that thrips population ranged from 0.32 to 0.35/5 leaves with pooled data 0.57 /5 leaves, aphid population varied from 1.05 to 1.95/5 shoots with pooled data 1.20/ five no of shoot, white fly from 4.03 to 5.41/5 leaves with pooled data 2.17 /5 leaves, yellow mites from 0.65 to 0.54/5 leaves with pooled data 0.77/5 leaves and fruit borer ranged from 0.21 to 0.40 /plant with pooled data 0.54 /plant though out the growth period. Similar results were also obtained by Nandini *et al.* (2010) [12] who conducted survey on capsicum pests under field condition at

Dharwad area. Sucking pests viz., thrips (0.60 to 2.30 thrips/leaves), mites (0.60 to 1.20 mites/3 leaves) and aphids (1.20 to 1.60 aphids/ 3leaves) were observed during vegetative to reproductive stage.

The pooled data obtained from yellow mite population gave an idea about opposite trend with the other insect like thrips, aphid, white fly and fruit borer. The population of yellow mite was positively correlated with the date of sowing. 1st December planted crop differ significantly from other two date but 1st January and 1st February were statistically at par. The findings were confirmed with the results obtained by Kethran *et al.* (2014) [13] conducted an experiment on effect on different sowing date on insect pest of chilli at Kunri, Mirpur Khas Sindh during 2014. He observed that late sowing (January 15th or January 30th) resulted in lower incidence of aphids, thrips, whitefly and fruit borer except mites. Such low level of insect pest caused a less crop injury which resulted in enhancing the green pod yield of chilli. Anon (2004) [14] also showed that early planting, as it is known in many crops, attracts greater intensity of pests and subsequent plant damage.

The percentage plant damage due to mite, percentage incidence of leaf curl virus, capsicum chlorosis virus and the marketable production of semi ripe green fruits are presented in table 3. The crops when planted in 1st February (18.3 t/ha) significantly increased the marketable production of fruits as compared to the 1st January (15.2 t/ha) and 1st December (13.3 t/ha) planted crops. It is justified by the findings of Kumar

(1995) [15]. He told that the yield losses range from 50-90 per cent due to insect pests of chilli. The damage due to mites and thrips together had been estimated to the tune of 50 per cent [16] and fruit borers is to an extent of 90 per cent [17].

In all treatments the percentage plant damage due to mite showed that, the 1st December planted crop (15.59%) was the best performing than the others two date of planting (21.77% and 18.45%) but the percentage incidence of leaf curl virus was low in 1st February (23.76%) than the 1st January (26.70%) and 1st December (34.58%) planted crops. The present study is justified with the conclusion by Bugti (2016) [18] who told that the highest pest population of Whitefly was recorded in capsicum spp. followed by Jassid, Thrips, Aphid, Mealy bug & Fruit borer. However, Venkatesh *et al.* (1998) [19] reported that chilli leaf curl complex was caused by leaf curl geminivirus (CLCV) transmitted by chilli mite (*Polyphagotarsonemus latus*), whitefly (*Bemisia tabaci*) and thrips (*Scirtothrips dorsalis*). The number of *B. tabaci* transmitted plant viruses has escalated and the total yield loss of important food and industrial crops has also increased [20]. The table 4 represented that the damage score in a different growth stages of plants in different dates of planting. In three dates of planting viz 1st December, 1st January and 1st February the highest damage score was obtained in 120th DAP (1.35 to 1.46) and lowest obtained in 60th DAT (0.68 to 0.85). The present findings confirmed with the results obtained by Alatawi *et al.* (2007) [21] who reported that older plants exhibited greater damage than younger plants.

Table 2: Population of bell pepper (var. Arka Mohini) pests on different dates of planting (Rabi, 2020-21 and 2021-22).

Date of planting	Thrips /5 leaves			Aphid/5 shoots (10 cm terminal portion)			White fly/5 leaves			Yellow mites/ 5 leaves			Larvae/plant		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
1 st December	1.81 (1.34)	1.96 (1.40)	1.24	3.46 (1.86)	5.45 (2.34)	2.09	5.67 (2.38)	7.65 (2.77)	2.57	0.15 (0.39)	0.22 (0.47)	0.42	0.91 (0.95)	1.01 (1.00)	0.97
1 st January	0.67 (0.82)	0.51 (0.72)	0.76	1.36 (1.17)	2.96 (1.72)	1.44	4.70 (2.17)	6.23 (2.50)	2.33	0.57 (0.76)	0.45 (0.67)	0.71	0.4 (0.67)	0.63 (0.80)	0.73
1 st February	0.32 (0.57)	0.35 (0.59)	0.57	1.05 (1.02)	1.95 (1.40)	1.20	4.03 (2.01)	5.41 (2.33)	2.17	0.65 (0.80)	0.54 (0.73)	0.77	0.21 (0.46)	0.40 (0.63)	0.54
C.D.(0.05)	0.06	0.04		0.08	0.10		0.09	0.07		0.06	0.04		0.04	0.04	

*Values in parenthesis are square root transformed.

Table 3: Percent plant damage and the marketable production of bell pepper (var. Arka Mohini) fruits in different dates of planting (Rabi, 2020-21 and 2021-22)

Date of planting	% damage of plant due to mite			% incidence of leaf curl virus			% incidence of capsicum chlorosis virus			Marketable Production(t/ha)		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Average
1 st December	7.86(16.28)	7.14(15.45)	15.59	25.71(30.46)	39.29(38.81)	34.58	14.14(22.79)	9.57(16.02)	18.67	14.25	12.42	13.34
1 st January	12.14(20.39)	8.57(17.02)	18.45	12.86(21.01)	29.29(32.76)	26.70	8.74(15.28)	9.41(16.75)	14.92	16.20	14.17	15.19
1 st February	15.71(23.57)	12.14(20.39)	21.77	10.71(19.09)	22.86(28.56)	23.76	3.76(10.32)	3.97(10.67)	7.54	19.63	16.95	18.29
C.D. (0.05)	4.20	3.02		4.28	5.45					30.02	15.44	

*Values in parenthesis are angular transformed (% data only)

Table 4: Damage score of bell pepper (var. Arka Mohini) plant in different dates of planting (Rabi, 2020-21 and 2021-22).

Days after planting	Damage score at 1 st December planting			Damage score at 1 st January planting			Damage score at 1 st February planting		
	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled	2020-21	2021-22	Pooled
60 th day	0.73	0.97	0.85	0.60	0.83	0.71	0.51	0.85	0.68
90 th day	1.40	1.41	1.40	1.19	1.20	1.20	1.15	1.26	1.21
120 th day	1.48	1.36	1.42	1.51	1.41	1.46	1.59	1.11	1.35
C.D.	0.21	0.28		0.24	0.20		0.17	0.19	
SE(m)	0.07	0.09		0.08	0.06		0.06	0.06	

Conclusion

It is inferred that 1st February planted crop reveals in lower incidence of aphids, thrips, whitefly and fruit borer and chlorosis virus, leaf curl virus except mites. Such low level of

insect pest caused a less crop injury which resulted in enhancing the yield of bell pepper. Pest and leaf curl virus incidence was moderate on 1st January planted crop.

References

1. Sreedhara DS, Kerutagi MG, Basavaraja H, Kunnal LB, Dodamani MT. Economics of capsicum production under protected conditions in Northern Karnataka. *Karnataka J. agric. Sci.* 2013;26(2):217-219.
2. Anonymous. www.indiastat.com. *Environ.* 2012;4:143-144.
3. Sunitha TR. Insect pests of *Capsicum annuum* var. *frutescens* (L.) and their management. M.Sc. (Entomology) thesis, University of Agricultural Sciences, Dharwad, Karnataka, India; c2007. p. 67.
4. Kaur S, Singh S. Efficacy of some insecticides and botanicals against sucking pests on capsicum under net house. *Agriculture for Sustainable Development.* 2013;1(1):25-29.
5. Ahmed M. Marketing of chillies: problems and prospects. Agriculture Marketing Information Service Directorate of Agriculture (Economics and marketing) Punjab Lahore, Pakistan; c2005.
6. Kumar M, Verma V. Bell Pepper (*Capsicum annuum* L.) production in low cost naturally ventilated poly houses during winters in the mid hills of India. *Acta Horticulture.* 2009;4:807.
7. Costa HS, Brown JK. Variation in biological characteristics and in esterase patterns among populations of *Bemisia tabaci* (Gennadius) and the association of one population with silver leaf symptom development. *Entomologia Experimentalis et Applicata.* 1991;61(3):211-219
8. Cohen S, Duffus JE, Liu HY. A new *Bemisia tabaci* biotype in the southwestern United States and its role in silver leaf of squash and transmission of lettuce infectious yellow virus. *Phytopathology.* 1992;82(1):86-90.
9. Niles GA. Breeding cotton for resistance to insect pests. In: *Breeding Plants Resistant to Insect.* Edited by Maxwell FG, Jennings P. Wiley J, Sons, New York; c1980. p. 657
10. Pathipati VL. Survey on insect pests, bioefficacy and dissipation studies of certain Insecticide molecules on capsicum (*Capsicum annuum* L. var. *grossum* Sendt.). PhD thesis submitted to Professor Jayashankar Telangana State Agricultural University; c2015.
11. Meena RS, Ameta OP, Meena BL. Population dynamics of sucking pests and their correlation with weather parameters in chilli, *Capsicum annum* L. *Crop. The Bioscan.* 2013;8(1):177-180.
12. Nandini Giraddi RS, Mantur SM, Mallapur CP. Survey and management of pests of capsicum under protected cultivation. Thesis submitted to University of Agricultural Sciences, Dharwad, Karnataka, India; c2010
13. Kethran MR, Sun YY, Khan S, Baloch SU, Wu LL, Yang TT, *et al.* Effect of different sowing date on insect pest population of chillies (*Capsicum annuum* L.). *Journal of Biology, Agriculture and Healthcare.* 2014;4(25):196-214.
14. Kumar NKK. Yield loss in chilli and sweet pepper due to *Scirtothrips dorsalis* Hood. (Thysanoptera: Thripidae). *Pest Management in Horticultural Ecosystem.* 1995;1(2):61-69.
15. Hosamani A. Management of chilli murda complex in irrigated ecosystem. Ph.D. Thesis. University of Agricultural Sciences. *Journal of Economic Entomology.* 2007;100:1904-1909.
16. Reddy MRS, Reddy GS. An eco-friendly method to combat *Helicoverpa armigera* (Hub.). *Insect Environ.* 1999;4:143-144.
17. Bugti GA. Varietal Preference of Insect Pests on Tomato Crop in District Naseerabad Balochistan Pakistan. *Journal of Entomology and Zoology Studies.* 2016;4(4):328-330
18. Venkatesh KM, Munniyappa V, Ravi KS, Krisnaprasad PR. Management of Chilli Leaf Curl complex. In: *Advances in IPM for Horticulture Crops,* Reddy PP, Kumar NKK, Verghese A(Eds.). Department of Agricultural Entomology, Tamil Nadu Agricultural University, Bangalore; c1998. p. 111-117.
19. Oliveira MRV, Henneberry TJ, Anderson P. History, current status and collaborative research projects for *Bemisia tabaci*. *Crop Protection.* 2001;20(9):709-723.
20. Alatawi FJ, Margolies DC, Nechols JR. Aesthetic damage thresholds for two spotted spider mites (Acari: Tetranychidae) on Impatiens: Effect of plant age and level of infestation. *Journal of Economic Entomology.* 2007;100(6):1904-1906.