



ISSN (E): 2277-7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2022; 11(12): 5118-5121
© 2022 TPI
www.thepharmajournal.com
Received: 14-09-2022
Accepted: 17-10-2022

DR Chaudhary
P.G. Student, Department of
Agronomy, COA, NAU,
Bharuch, Gujarat, India

Vaishali H Surve
Assistant Professor, Department
of Agronomy, COA, NAU,
Bharuch, Gujarat, India

SM Bambhaneeya
Assistant Professor, Department
of Soil Science and Agril.
Chemistry, COA, NAU,
Bharuch, Gujarat, India

AD Raj
Associate Professor, Department
of Agronomy, COA, NAU,
Bharuch, Gujarat, India

SP Deshmukh
Assistant Research Scientist
(Agronomy) Niger Research
Station, NAU, Varanasi,
Gujarat, India

Corresponding Author:
DR Chaudhary
P.G. Student, Department of
Agronomy, COA, NAU,
Bharuch, Gujarat, India

Effect of organics on yield, quality, nutrient content and uptake of soybean [*Glycine max* (L.)] under rainfed condition

DR Chaudhary, Vaishali H Surve, SM Bambhaneeya, AD Raj and SP Deshmukh

Abstract

A field experiment was conducted during 2021 to study the effect of organics as foliar soil application on yield, quality, nutrient content and uptake of soybean [*Glycine max* (L.)] under rainfed condition. The results revealed that foliar application of 4% seaweed extract at 30 and 45 DAS recorded significantly higher seed yield (1595 kg ha⁻¹), stover yield (3062 kg ha⁻¹), crude protein yield (634 kg ha⁻¹), oil yield (312 kg ha⁻¹), nutrient content and uptake by seed and stover of soybean. Soybean sown with application of jeevamrut @ 500 l ha⁻¹ produced significantly the highest seed yield (1644 kg ha⁻¹), stover yield (3072 kg ha⁻¹), crude protein yield (653 kg ha⁻¹), oil yield (319 kg ha⁻¹), nutrient content and uptake by seed and stover of soybean. The interaction effect of 4% seaweed extract at 30 and 45 DAS and jeevamrut @ 500 l ha⁻¹ was recorded significantly higher seed yield of soybean.

Keywords: Soybean, organic spray, jeevamrut, yield, quality, nutrient, content and uptake

Introduction

Soybean [*Glycine max* (L.)] is an important oil yielding crop having worldwide adaptation. It belongs to the legume family and native to East Asia. It belongs to the family Fabaceae and sub family Papilionaceae. Soybean is known as the “Golden Bean” of the twentieth century. The leading soybean growing countries are USA, Brazil, China, Argentina, and India. In India, soybean occupies an area of about 10.38 million hectares producing 10.94 million tonnes with the productivity of 1050 kg ha⁻¹, whereas in Gujarat it is grown in 0.15 million hectares with the production of 0.12 million tonnes and productivity of 788 kg ha⁻¹ (Anonymous, 2020) [2]. Panchgavya is an organic product blended with five different cow products, commonly applied to crop plant in organic farming. It acts as growth promoter and immunity booster and also restricts the incidence of common disease. Vermiwash is watery extract of vermicompost, extracted in the presence of rich population of earthworm and contains several enzymes, plant growth hormone, vitamins along with micro and macro and macronutrients which increase the resistance power of crops against various diseases and enhance the growth and productivity of crops (Zambare *et al.* 2008) [11].

The extract of marine macro algae *viz.*, brown, red and green algae is known to have positive effect on growth and yield of crops. The brown algae are the most commonly used seaweeds in agriculture. Cattle urine is a good source of nitrogen, phosphate, potassium, calcium, magnesium, chlorite and sulphate. It contains 95% water, 2.5% urea, 2.5% others (mineral salts, hormones and enzyme). Novel organic liquid nutrient is a product of Navsari Agricultural University which was patented in the year of 2012. It is prepared from banana pseudostem sap. Jeevamrut also contains enormous amount of microbial load which multiply and enhance N fixation and nutrient mobilization and utilization increase the soil fertility (Palekar, 2006) [5]. Jeevamrut contains small amount of macro and micronutrient as well as growth hormone which is helpful to enhancing the growth and yield of crops. Looking into importance of organics as foliar spray and soil application of jeevamrut, a field experiment was conducted to see that effects on growth, yield and economics of soybean under rainfed condition.

Materials and Methods

The field experiment was conducted at the College Farm, Navsari Agricultural University, campus Bharuch (South Gujarat Agro Climatic Zone - II) in the Plot No. 12 during the *Kharif*

season of 2021. The soil of experimental field was clayey in texture and slightly alkaline in reaction. The soil was low in available N (244 kg ha⁻¹), low in available P₂O₅ (26.2 kg ha⁻¹) and high in available K₂O (330.21 kg ha⁻¹). The experiment was laid out in FRBD with 12 treatment combinations consisting of two factors *i.e.*, foliar nutrition application (F₁-control, F₂- 4% panchgavya at 30 and 45 DAS, F₃- 2% novel organic liquid at 30 and 45 DAS, F₄-4% seaweed extract at 30 and 45 DAS, F₅- 4% cow urine at 30 and 45 DAS and F₆-2% vermiwash at 30 and 45 DAS) and soil application (S₁-control, S₂-jeevamrut @ 500 l ha⁻¹) were tested in factorial randomized block design with three replication. The entire dose of fertilizer was 30-60-00 N: P₂O₅: K₂O kg ha⁻¹. Soybean variety KDS-344 was sown by the opening of furrow at distance of 45 x 10 cm. Specially prepared jeevamrut solution was applied at the rate of 500 l ha⁻¹ along with the application of FYM in treated plots. Applications of different foliar spray as per treatments were applied at 30 and 45 DAS, using 500-liter water hectare⁻¹.

The data on seed and stover yield was recorded from the net plot and converted on a hectare basis. The nitrogen content in soybean seed was estimated by micro alkaline permanganate oxidation method as described by Subbiah and Asija (1956)^[9]. The crude protein content of the seed was computed by multiplying the nitrogen percentage with 6.25 for each treatment. The oil content was determined by Soxhlet extraction method as per procedure given by Soxhlet (1879)^[13]. Chemical studies about nitrogen, phosphorus, potassium content and their uptake by seed and stover and available nitrogen, phosphorus, potassium status in the soil after harvest of the crop were determined as per different methods *viz.*, Modified alkaline permanganate oxidation method (for N), Spectro photometric (for P) and Flame photometric method (for K). The data were analyzed statistically by adopting the standard procedures described by Panse and Sukhatme (1967)^[6]. The purpose of the analysis of variance was to determine the significant effect of treatments on soybean. Uptake of nutrients by seed and plant was calculated by using following formula:

$$\text{Nutrient uptake (kg ha}^{-1}\text{)} = \frac{\text{Nutrient content (\%)} \times \text{seed / stover yield (kg ha}^{-1}\text{)}}{100}$$

Results and Discussion

Effect of foliar nutrition

Yield

The data given in Table 1 indicated that significantly higher seed yield (1595 kg ha⁻¹) and stover yield (3062 kg ha⁻¹) was registered under foliar application of 4% seaweed extract at 30 and 45 DAS (F₄), however it was statistically at par with treatment of 4% panchgavya at 30 and 45 DAS (F₂), 2% novel organic liquid at 30 and 45 DAS (F₃) and 2% vermiwash at 30 and 45 DAS (F₆). Higher seed yield might be due to IAA and GA present in seaweed extract when spray could have created stimuli in the plant system which in turn increased the production of growth regulators in cell system and the action of growth regulators in plant system stimulated the necessary growth and development coupled with better translocation and accumulation of photosynthates from source to sink and ultimately increased the seed yield. These findings were in accordance with the results of Akhila *et al.* (2017)^[11], in their studies foliar application of 2% *Kappaphycus alvarezii*

seaweed sap gave significantly the highest seed (26.65 q ha⁻¹) and stover yield (12.21 q ha⁻¹) over 6% *Kappaphycus alvarezii*, 1% enriched banana pseudostem sap and 2% enriched banana pseudostem sap.

Quality parameters

The data of Table 1 indicated that protein content and oil content were not differed significantly due to different foliar nutrition practices. Numerically higher crude protein content (39.77 per cent) and oil content (19.61 per cent) were recorded by the application of 4% seaweed extract at 30 and 45 DAS (F₄). However, protein yield and oil yield were significantly influenced due to different foliar nutrition practices. Application of 4% seaweed extract at 30 and 45 DAS (F₄) recorded significantly higher protein yield (634 kg ha⁻¹) and oil yield (312 kg ha⁻¹) which was statistically at par with treatment of 4% panchgavya at 30 and 45 DAS (F₂), 2% novel organic liquid at 30 and 45 DAS (F₃) and 2% vermiwash at 30 and 45 DAS (F₆). This might be due to increase in seed yield when applied with seaweed extract. Similar reasons have also been reported by Zodape *et al.* (2010)^[12]. They reported that protein content had increased by 7.07% with the foliar application of 10% *Kappaphycus alvarezii* seaweed sap as compared to 5% and 15%.

Nutrient content and uptake

An appraisal of data given in Table 3 revealed that no significant difference was observed in nitrogen, phosphorus and potassium content in seed and stover due to different foliar nutrition practices. However, significantly higher nitrogen, phosphorus and potassium uptake by seed and stover was recorded by application of 4% seaweed extract at 30 and 45 DAS (F₄) which was remained statistically at par with treatment of 4% panchgavya at 30 and 45 DAS (F₂), 2% novel organic liquid at 30 and 45 DAS (F₃) and 2% vermiwash at 30 and 45 DAS (F₆) and lowest nitrogen, phosphorus and potassium uptake by seed and stover was recorded in control (F₁). This increased might be due to foliar application of seaweed extract. The presence of marine bioactive substance in seaweed extract improves stomata uptake efficiency in treated plants as compared to non-treated plants. Similar finding by Pramanick *et al.* (2013) recorded highest N (57.67 kg ha⁻¹), P (5.85 kg ha⁻¹) and K (22.97 kg ha⁻¹) uptake by seed and highest N (143.33 kg ha⁻¹), P (7.83 kg ha⁻¹) and K (80.95 kg ha⁻¹) uptake by stover by the application of 15% *Kappaphycus*-sap + RDF as compared to different concentration of *Kappaphycus*-sap and *Gracilaria*-sap.

Available nutrient in soil after harvest

The different foliar nutrition practices were influenced non-significant effect on available N, available P₂O₅ and K₂O in the soil after harvest of soybean crop. Dwivedi *et al.* (2015) recorded maximum available N (166.2 kg ha⁻¹), P₂O₅ (34.05 kg ha⁻¹) and K₂O (418.9 kg ha⁻¹) in soil after harvest under foliar spray of 15% *Kappaphycus* spp + RDF on soybean.

Effect of soil application

Yield

The data given in Table 1 indicated that significantly the highest seed yield (1644 kg ha⁻¹) and stover yield (3072 kg ha⁻¹) of soybean was observed under the application of jeevamrut @ 500 l ha⁻¹ (S₂) in comparison to control (S₁) (1290 kg ha⁻¹, 2466 kg ha⁻¹). The significant improvement in seed yield

plant¹ was attributed due to increased supply of plant nutrients, more root nodules and phytohormones supplied through application of jeevamrut. The results were supported by findings of Upperi *et al.* (2009) [10], who recorded application of jeevamrut recorded the highest mean yield and yield attributes of treated field as compared to control.

Quality parameters

The data of Table 1 indicated that protein content and oil content were not differed significantly due to soil application. Numerically highest crude protein content (39.69 per cent) and oil content (19.39 per cent) were recorded by application of jeevamrut @ 500 l ha⁻¹ (S₂) in comparison to control (S₁). However, protein yield and oil yield were significantly influenced due to soil application. Jeevamrut @ 500 l ha⁻¹ (S₂) recorded significantly highest protein yield (653 kg ha⁻¹) and oil yield (319 kg ha⁻¹) in comparison to control (S₁). Increase in protein yield might be due to increase in nitrogen content and seed yield by soil application of jeevamrut. Similar finding reported by Patil and Udmale (2016) [7] for the higher protein and oil content in seed of soybean by the application of FYM + vermicompost (50% each) + jeevamrut (2 times *i.e.*, at 30 and 45 DAS).

Nutrient content and uptake

An appraisal of data given in Table 3 revealed that no significant difference was observed in nitrogen, phosphorus and potassium content in seed and stover due to soil application of jeevamrut @ 500 l ha⁻¹. However, significantly highest nitrogen, phosphorus and potassium uptake by seed and stover was recorded by application of jeevamrut @ 500 l ha⁻¹ (S₂) and lowest nitrogen, phosphorus and potassium uptake by seed and stover was recorded in control (S₁). This is might be due to quick build-up of soil micro flora and fauna which has consequently increased the enzymatic activity helped in mineralization, solubilization of native and applied

nutrients and making them available for plant uptake. These results are in accordance with the results of Patil and Udmale (2016) [7], who reported that application of FYM + vermicompost (50% each) + jeevamrut (2 times *i.e.*, at 30 and 45 DAS) gave significantly higher uptake of N, P and K by soybean.

Available nutrient in soil after harvest

The soil application was influenced non-significant effect on available N, available P₂O₅ and K₂O in the soil after harvest of soybean crop. Lahariya *et al.* (2013) [14] recorded the maximum available N, P, K content in soil after harvesting of soybean as compared to control when application of jeevamrut @ 500 l ha⁻¹.

Interaction effect

The data in Table 3 revealed that the treatment combination of foliar nutrition (F) and soil application (S) significantly influenced seed yield of soybean. The treatments combination of 4% seaweed extract at 30 and 45 DAS with jeevamrut @ 500 l ha⁻¹ gave significantly higher seed yield (1745 kg ha⁻¹) and were found statistically at par with application of 2% vermivash at 30 and 45 DAS with *jeevamrut* @ 500 lit ha⁻¹ (1725 kg ha⁻¹), application of 4% *panchgavya* at 30 and 45 DAS with *jeevamrut* @ 500 lit ha⁻¹ (1685 kg ha⁻¹) and application of 2% Novel organic liquid at 30 and 45 DAS with *jeevamrut* @ 500 lit ha⁻¹ (1619 kg ha⁻¹) while lowest number seed yield (936 kg ha⁻¹) was registered with treatments combination of control. This might be due to the combine beneficial effect of seaweed extracts and jeevamrut which contain nutrients, plant growth hormones auxins, cytokinin and gibberellins as well as other plant bio stimulant like amino acid and which is increase the photosynthetic activity and further, the translocation and accumulation of photosynthates.

Table 1: Effect foliar nutrition and soil application on yield and quality of soybean

Treatments	Yield (kg ha ⁻¹)		Crude protein content (%)	Crude protein yield (kg ha ⁻¹)	Oil content (%)	Oil yield (kg ha ⁻¹)
	Seed	Stover				
Foliar nutrition (F)						
F ₁	1238	2428	37.63	470	18.20	226
F ₂	1562	2873	39.63	618	19.40	303
F ₃	1526	2850	39.53	603	18.97	290
F ₄	1595	3062	39.77	634	19.61	312
F ₅	1374	2602	38.94	538	18.64	257
F ₆	1505	2797	39.23	592	19.04	287
S Em ±	44.96	136.30	0.67	20.81	0.48	11.54
C D at 5%	131.87	399.75	-	61.03	-	33.87
Soil application (S)						
S ₁	1290	2466	38.55	499	18.57	240
S ₂	1644	3072	39.69	653	19.39	319
S Em ±	25.96	78.69	0.39	12.01	0.27	6.66
C D at 5%	76.13	230.79	-	35.23	-	19.55
Interaction (F x S)						
S Em ±	63.58	192.75	0.96	29.43	0.68	16.33
C D at 5%	186.49	-	-	-	-	-

Table 2: Seed yield of soybean as influenced by interaction effect of foliar nutrition and soil application.

Treatments	Soil application (S)	
	S ₁	S ₂
F ₁	936	1542
F ₂	1440	1685
F ₃	1434	1619
F ₄	1446	1745
F ₅	1199	1551
F ₆	1285	1725
S Em ±	63.58	
CD at 5%	186.49	

Table 3: Nutrient content, uptake and available nutrient status of soil by soybean as influenced by foliar nutrition and soil application

	Nutrient content (%)						Nutrient uptake (kg ha ⁻¹)						Available Nutrients (kg ha ⁻¹)			
	Seed			Stover			Seed			Stover			N	P ₂ O ₅	K ₂ O	
	N	P	K	N	P	K	N	P	K	N	P	K				
Foliar nutrition (F)																
F ₁	6.021	0.354	1.233	1.245	0.217	0.810	75.23	4.39	15.31	30.30	5.28	19.63	240.61	24.322	291.39	
F ₂	6.340	0.369	1.271	1.297	0.233	0.852	98.96	5.78	19.87	37.47	6.67	24.62	246.70	26.538	301.41	
F ₃	6.325	0.361	1.251	1.292	0.228	0.840	96.58	5.52	19.16	36.58	6.51	23.96	246.02	25.893	298.13	
F ₄	6.362	0.372	1.276	1.311	0.238	0.873	101.54	5.95	20.38	40.32	7.31	26.84	248.69	26.698	304.29	
F ₅	6.230	0.356	1.241	1.271	0.226	0.828	86.09	4.91	17.12	33.13	5.90	21.57	244.48	25.506	295.51	
F ₆	6.276	0.360	1.249	1.287	0.227	0.830	94.76	5.44	18.85	36.04	6.37	23.15	245.38	25.727	296.64	
S Em. ±	0.109	0.010	0.028	0.031	0.007	0.020	3.32	0.21	0.74	1.86	0.36	1.15	5.481	0.96	6.49	
CD at 5%	-	-	-	-	-	-	9.76	0.63	2.19	5.46	1.06	3.39	-	-	-	
Soil application (S)																
S ₁	6.168	0.354	1.231	1.258	0.224	0.822	79.90	4.57	15.91	31.03	5.54	20.29	242.40	25.023	292.67	
S ₂	6.350	0.370	1.275	1.310	0.232	0.855	104.49	6.09	20.99	40.25	7.14	26.30	248.03	26.538	303.12	
S Em. ±	0.063	0.006	0.016	0.018	0.004	0.012	1.92	0.12	0.43	1.07	0.20	0.66	3.164	0.55	3.75	
CD at 5%	-	-	-	-	-	-	5.63	0.36	1.26	3.15	0.61	1.96	-	-	-	
Interaction (F x S)																
S Em. ±	0.154	0.014	0.040	0.043	0.010	0.029	4.70	0.30	1.05	2.63	0.51	1.63	7.751	1.35	9.18	
CD at 5%	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

Conclusions

On the basis of experiment, it can be concluded that soil application of jeevamrut @ 500 lit ha⁻¹ prior to sowing of soybean with foliar application of 4% seaweed extract or 4% panchgavya or 2% novel organic liquid or 2% vermiwash at 30 and 45 DAS found better quality, yield and nutrient uptake of soybean and maintaining soil health.

References

- Akhila K, Kaswala AR, Priyanka, Dubey PK. Effect of liquid fertilizers on growth yield and economics of the green gram (*Vigna radiata*) crop under organic farming. Int. J Chem. Stud. 2017;5(6):809-812.
- Anonymous. Ministry of Agriculture and FW, GOI. Agriculture Statistics at a Glance Archives, 2020;20(2):1672-1674.
- Dwivedi SK, Ashok PK. Impact of natural organic fertilizer (seaweed saps) on productivity, nutrient uptake and economics of greengram (*Phaseolus radiata* L.). Int. J Environ. Sci. 2015;9(4):461-466.
- Laharia GS, Patil DU, Damre PR. Effect of organic sources on soil fertility, nutrient uptake and yield of soybean. Crop Res. 2013;45(1-3):155-159.
- Palekar S. Text book on shoonya bandovalada naisargika krushi, published by swamy Anand. Agri Prakashana, Bangalore, 2006, pp. 67.
- Panse VG, Sukhatme PV. Statistical Methods for Agricultural Workers. ICAR Publication, Ministry of Agriculture, New Delhi, 1973.
- Patil HM, Udmale KB. Response of different organic inputs on growth and yield of Soybean on *Inceptisol*. Schor. J Agri. Sci. 2016;6(5):139-144.
- Pramanick B, Brahmachari K, Ghosh A. Effect of seaweed saps on growth and yield improvement of greengram. Afri. J Agri. Res. 2013;8(13):1180-1186.
- Subbiah BV, Asija GL. A rapid procedure for the estimation of available nitrogen in soil. Curr. Sci. 1956;25:259.
- Upperi SN, Lokesh BK, Maraddi GN, Agnal MB. Jivamrut, a new organic approach for disease management and crop production in groundnut. Environ. and Eco. 2009;27(1):202-204.
- Zambare VP, Padul MV, Yadav AA, Shete TB. Vermiwash: biochemical and microbiological approach as ecofriendly soil conditioner. J agric, Bio. Sci. 2008;3(4):1-5.
- Zodape ST, Mukhopadhyay K, Eshwaran, Reddy MP, Chakara J. Enhancement of yield and nutritional quality in greengram treated with seaweed *Kappaphycus alvarezii*. J Sci. and Indu. Res. 2010;69(1):468-471.
- Soxhlet F. Dinglers Polytech. J. 1879;232:461-465.
- Lahariya C, Subramanya BP, Sosler S. An assessment of hepatitis B vaccine introduction in India: Lessons for roll out and scale up of new vaccines in immunization programs. Indian journal of public health. 2013 Jan 1;57(1):8.