



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2022; 11(4): 1669-1676
© 2022 TPI

www.thepharmajournal.com

Received: 23-02-2022

Accepted: 31-03-2022

Priyavart Mishra

Department of Soil Science and
Agricultural Chemistry, C.S.
Azad University of Agriculture
and Technology, Kanpur,
Uttar Pradesh, India

Sanjeev Sharma

Department of Soil Science and
Agricultural Chemistry, C.S.
Azad University of Agriculture
and Technology, Kanpur,
Uttar Pradesh, India

US Tiwari

Department of Soil Science and
Agricultural Chemistry, C.S.
Azad University of Agriculture
and Technology, Kanpur,
Uttar Pradesh, India

Hanuman Prasad Pandey

Department of Soil Science and
Agricultural Chemistry, C.S.
Azad University of Agriculture
and Technology, Kanpur,
Uttar Pradesh, India

RK Pathak

Department of Soil Science and
Agricultural Chemistry, C.S.
Azad University of Agriculture
and Technology, Kanpur,
Uttar Pradesh, India

AK Sachan

Department of Soil Science and
Agricultural Chemistry, C.S.
Azad University of Agriculture
and Technology, Kanpur,
Uttar Pradesh, India

Corresponding Author:

Sanjeev Sharma

Department of Soil Science and
Agricultural Chemistry, C.S.
Azad University of Agriculture
and Technology, Kanpur,
Uttar Pradesh, India

Response of INM on nutrient concentration and their uptake of maize (*Zea mays* L.) crop

Priyavart Mishra, Sanjeev Sharma, US Tiwari, Hanuman Prasad Pandey, RK Pathak and AK Sachan

Abstract

A field experiment was conducted at field no. 6 Student's In structural Farm at Chandra Shekhar Azad University of Agriculture and Technology, Kanpur during the *Kharif* season 2017 to find out integrated nutrient management effect on maize with ten treatments i.e. T₁ (125% RDN), T₂ (100% RDN), T₃ (100% RDN + 25% N FYM), T₄ (100% RDN + 25% N FYM + S₃₀), T₅ (100% RDN + 25% N FYM + S₃₀ + Zn₅), T₆ (75% RDN), T₇ (75% RDN + 25% N FYM), T₈ (75% RDN + 25% N FYM + S₃₀), T₉ (75% RDN + FYM + S₃₀ + Zn₅), T₁₀ (Control) in RBD with 3 replications. Maize variety Azad Uttam was taken for study. The results revealed that the N, P, K, S and Zn content in maize grain varies from 1.12 to 1.44%, 0.26-0.37%, 0.32-0.44%, 0.28-0.39% and 22.20-31.80% mg/kg and the N, P, K, S and Zn content of maize stalk varies from 0.34 to 0.55%, 0.110 to 0.142, 1.02 to 1.19%, 0.20 to 0.33% and 14.20-22.80 mg/kg respectively. The total uptake values of N, P, K, S and Zn varied from 43.27 to 107.58 kg/ha, 11.47 to 26.95 kg/ha, 67.19 to 132.11 kg/ha, 17.24 to 46.07 kg/ha and 1268.24 to 3355.10 g/ha. The maximum nutrient concentration and uptakes in case of all treatments was found in T₅ (100% RDN + 25% N FYM + S₃₀ + Zn₅) and lowest in T₁₀ (Control).

Keywords: *Zea mays*, nutrient concentration, uptake, Azad Uttam, FYM, grain, stalk

Introduction

Maize (*Zea mays* L.) is one of the most important cereal crop, next to rice and wheat and is used as a food for human and feed for animals. This crop has been developed into a multi dollar business in countries viz. Thailand, Taiwan, Singapore, Malaysia, USA, Canada and Germany, because of its potential as a value added product for export and a good food substitute. In India maize is primarily a *kharif* crop which is sown just before monsoon starts. This crop usually grows well under temperatures varying from 22°C to 30°C, although it can tolerate temperatures as high 35°C. This crop is affected by frost, so it is grown where at least 5 frost free months are available in the year. It requires at least 50 to 90 mm of rainfall. It is not recommended to cultivate this crop in the areas where rainfall is more than 100 mm. Maize or corn can be cultivated successfully in variety of soils ranging from clay loam to sandy loam to black cotton soil. For better yield of maize, soils should consider good organic matter content having high water holding capacity. Well drained soils with pH of 5.5 to 7.0 are preferred for maize farming. Maize is grown in an area of 9.76 million hectares with production of 26.14 million tonnes and productivity of 2629.28 kg ha⁻¹ (Government of India, 2017). Maize yield is generally higher in high solar intensities, lower night temperature and lower pest infestation. Optimum plant density leads to better utilization of solar radiation resulting into corn dry matter accumulation and biomass production. Uttar Pradesh is the major producing state contributes 60 percent area and 70 percent of maize production in India. Kumar (2008) [4] reported that application of 40, 80 and 120 kg N ha⁻¹ tended to increase nitrogen uptake by pop corn by 46.5, 78 and 99 percent, respectively over control. Kumar and Singh (2010) [3] found the DTPA-extractable Zn which was higher in the treatment combination of 100% NPK + 10 kg Zn ha⁻¹. They concluded that integrated application of 100% NPK + green gram + 5.0 t FYM was the most effective treatment in meeting the micronutrient requirements of rice crop.

Kumar *et al.* (2007) [2] reported that application of 100 percent RDF recorded significantly higher dry matter production in leaf, stem, cob, yield and yield attributes and uptake of the nitrogen, phosphorus, potassium and also higher net returns (Rs.20,898 ha⁻¹) and B:C ratio (2.14) over 50% RDF and it was found to be on par with 75% RDF.

Laxminarayana (2006) ^[5] recorded the maximum total uptake of nitrogen, phosphorus and potassium with the value of 73.27, 21.93 and 87.86 kg ha⁻¹, respectively by rice in treatment where 50% NPK along with 50% N through green manure was applied in *Kharif*.

Laxminarayana and Patiram (2006) ^[5] revealed that application of optimum doses of NPK in combination with green manure @ 5 t ha⁻¹ recorded highest uptake of N, P and K followed by 100 per cent NPK + poultry manure and 100 per cent NPK + FYM. However, the uptake responses were high with the balanced application of NPK in comparison to sub optimal and super optimal doses of NPK an Ultisol

Ramesh *et al.* (2008) ^[8] reported that chemical fertilizer treatment recorded significantly the highest total nitrogen and potassium uptake (332.2 and 343.6 kg ha⁻¹) which was on par with poultry manure application (294.1 and 319.7 kg ha⁻¹, respectively) in maize- linseed cropping system.

Roul and Sarawgi (2005) ^[9] revealed that recommended dose of N (RDN) blended with FYM and 100% RDN + 5 tons of FYM are better in respect of grain yield, straw yield and N-content in grain than other treatments on pooled data basis.

Setia and Sharma (2007) ^[11] reported that phosphorus uptake by grain (10.4 kg ha⁻¹) and stover (13.0 kg ha⁻¹) in maize was highest when N, P and K were applied @ 180:35:33 kg ha⁻¹ as compared to lower doses of fertilizers.

Sujatha *et al.* (2008) ^[13] reported that maximum uptake of N, P and K by maize crop to the extent of 241.1, 35.2 and 234.2 kg ha⁻¹, respectively was observed with application of sunhemp green manure + poultry manure + 100% RDN over control.

Thavaprakash and Velayudham, (2007) ^[14] conducted on the effect of integrated nutrient management practices on nutrient uptake revealed that the nutrient removal by maize under combined application of FYM, inorganic fertilizers and bio fertilizer as *Azospirillum* registered higher NPK uptake over (150 + 60 + 40 kg ha⁻¹ NPK) recommended dose of fertilizer.

Varalakshmi *et al.* (2005) ^[15] reported increase in organic carbon content of sandy clay loam soil from 8.1 g kg⁻¹ to 8.5 g kg⁻¹ after finger millet crop without addition of any fertilizers which was comparatively lower than integrated application of 50 per cent nitrogen through fertilizer and 50 per cent through FYM (9.2 g kg⁻¹).

Materials and Methods

The experiment was conducted on Maize during *kharif* season of 2017 under natural condition at field no. 6 Student's In structural Farm at Chandra Shekhar Azad University of Agriculture and Technology, Kanpur. The soil of the experimental field was alluvial in origin. Soil sample (0-15cm) depths were initially drawn from randomly selected parts of the field before sowing. The quantity of soil sample was reduced to about 500 gm through quartering technique. The soil sample was then subjected to mechanical and chemical analysis in order to determine the textural class and fertility status the soils were sampled to a depth of 0-30 cm of the soil, air-dried and sieved (2 mm) for soil analyses. Some physical and chemical properties of soils are given in Table 1.

Table 1: Some properties of the <2mm fraction of the top 30 cm of soil used for the site

S. No.	Particulars	Values
A.	Mechanical separates	
1.	Sand (%)	59.6
2.	Silt (%)	17.4
3.	Clay (%)	23.00
4.	Textural Class	Sandy loam
B.	Physico-chemical properties	
5.	pH (1:2.5)	8.2
6.	EC (1:2.5) (dS/m at 25°C)	0.20
7.	Organic Carbon (%)	0.36
8.	Available Nitrogen (kg/ha)	190.00
9.	Available Phosphorus (kg/ha)	13.50
10.	Available Potassium (kg/ha)	182
11.	Available Sulphur (kg/ha)	15.80
12.	Available Zinc (ppm)	0.56
13.	Particle Density (Mg/m ³)	2.54
14.	Bulk Density (Mg/m ³)	1.30
15.	Pore Space (%)	46.0

Maize variety Azad Uttam was taken for study. In the present experiment 10 treatments T₁ (125% RDN), T₂ (100% RDN), T₃ (100% RDN + 25% N FYM), T₄ (100% RDN + 25% N FYM + S₃₀), T₅ (100% RDN + 25% N FYM + S₃₀ + Zn₅), T₆ (75% RDN), T₇ (75% RDN + 25% N FYM), T₈ (75% RDN + 25% N FYM + S₃₀), T₉ (75% RDN + FYM + S₃₀ + Zn₅), T₁₀ (Control) were laid out in Randomized Block Design (RBD) with three replications having plot size 5 x 4 meter square. Doses of fertilizers are applied @ 120 Kg N, 60 Kg P₂O₅, 40 Kg K₂O/ha 30 Kg S/ha, 5 Kg Zn/ha and Organic manure 60 tonne/hathrough Urea, D.A.P and Murate of Potash, Elemental sulphur, Zinc oxide and Farm Yard Manure. Sowing is done @ 20 kg seed ha⁻¹ maize variety Azad Uttam was used and sown on 22 June 2017. Row to row and plant to plant distance remain 60 and 20 respectively. Seed were sown about 5-6 cm depth.

Field Preparation: The experimental field was ploughed once with soil turning plough followed by two cross harrowing. After each operation, planking was done to level the field and to obtain the fine tilth. Finally layout was done and plots were demarked with small sticks and rope with the help of manual labour in each block.

Application of fertilizers: The crop was fertilized as per treatment. The recommended dose of nutrient i.e. N, P, and K was applied @ 120: 60: 40 kg ha⁻¹ respectively.

Time and method of fertilizer: Half does N₂ and total phosphorus, potash, zinc and sulphur were applied as basal dressing. Remaining dose of nitrogen was applied through top dressing after knee-high stage. Well decompose FYM applied @ 60 t ha⁻¹ 15 day after sowing.

Seed Treatment: To ensure the seeds free from seed borne diseases, seeds were treated with thiram 75% WDP (1.5g/kg of seed).

Seed and sowing: 20 kg seed ha⁻¹ maize variety Azad Uttam was used and sown on 22 June 2017. Row to row and plant to plant distance remain 60 and 20 respectively. Seed were sown about 5-6 cm depth.

Intercultural operations: Weeding and hoeing were done with khurpi and hand hoe after germination.

Irrigation: Tube-well was the source of irrigation. Irrigation was provided in the crop as and when required.

Harvesting: The crop was harvested at proper stage of maturity as determined by visual observations. Half meter length on either end of each plot and two border rose from each side as border were first removed from the field to avoid error. The crop in net plot was harvested for calculation on yield data. Produce was tied in bundles and weighted for biomass yield. Threshing of produce of each net crop was done by manually.

Soil Analysis

Mechanical Separates: Soil separates analyzed by International pipette method as described by the Piper (1966). **pH:** pH of the soil determined by using soil water suspension (1:2.5) with the help of digital pH meter. **EC:** EC also determined using soil water suspension (1:2.5) with help of conductivity meter (Jackson, 1967) [3]. **Organic Carbon:** Organic Carbon was determined by Walkley and Black's rapid titration method as described by Jackson (1967) [3]. **Available Nitrogen:** It was determined by Alkaline Potassium Permanganate Method described by Subbiah and Asija (1956) [20]. **Available Phosphorus:** It is determined by Olsen's method using 0.5 M NaHCO₃ (Olsen *et al.* 1954) [12]. **Available Potassium:** Potassium is determined by using Neutral Normal Ammonium Acetate (pH 7.0) by Flame Photometer. **Available Sulphur:** Available Sulphur was determined by turbidimetric method (Chesnin and Yien, 1950) [1] after extraction with 0.15% CaCl₂ solution. **Available Zinc:** Available Zn is determined by Atomic Absorption Spectrophotometer with the help of DTPA extractant (Lindsey and Norvell, 1978) [9].

Plant Analysis

Plant samples were dried first in air then kept in oven at 70°C for 8 hr to make the sample free from excess moisture. The samples were grounded in a Wiley mill having stainless parts and stored in polythene bags.

Preparation of extract: Fine ground plant samples were digested in triacid mixture of conc. Nitric acid, sulphuric acid and perchloric acid for P and K determination in 10:4:1 ratio. Diacids (9:4 mixture of HNO₃ and HClO₄) digestion method is adopted for Zn extraction.

Determination of N, P, K, S and Zn in plant

- Nitrogen:** N is determined by Kjeldahl method given by Jackson (1967) [3].
- Phosphorus:** P is determined colorimetrically by vanadate-molybdate yellow colour method as advocated by Chapman and Pratt (1961) [14].
- Potassium:** K determination has been done using flame photometric method (Chapman and Pratt, 1961) [14] outlined by Jackson (1967) [3].
- Sulphur:** S is determined through turbidimetric method (Chesnin and Yien, 1956) [1].

- Zinc:** Zn is extracted from plant with the help of atomic absorption spectrophotometer (Lindasey and Norwell, 1978) [9].

Uptake

To calculate the uptake of N, P, K and S in grain as well as in straw, the following formula is used-

$$\text{Uptake of Nutrients (kg/ha)} = \frac{\text{Nutrient content(\%)} \times \text{Yield(kg/ha)}}{100}$$

Statistical Analysis: The data on various characters studied during the course of investigation were statistically analyzed for randomized block design. Wherever treatment differences were significant ("F" test), critical differences were worked out at five per cent probability level. The data obtained during the study were subjected to statistical analysis using the methods advocated by Chandel (1990) [2].

Results

Effect of INM on Nutrient Concentration

Nitrogen concentration

Grain

Data pertaining to N concentration in grain given in table 2 showed significant increase in all the treatment in comparison to its respective control. N concentration in grain varied from 1.22% (control) to 1.44% in T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹). Variation in N concentration within 75% RDN, 100% RDN and 125% RDN was noted significant. Integration of S, FYM and Zn influenced N concentration when applied with 75% RDN and 100% RDN treatment. It was also observed that integration of S showed higher increase in N concentration over Zn and FYM at 75% RDN and 100% RDN levels.

Table 2: Effect of Integrated Nutrient Management on N content (%) in grain and stalk

S. No.	Treatments	N content (%) in grain	N content (%) in stalk
1.	T ₁	1.34	0.46
2.	T ₂	1.30	0.42
3.	T ₃	1.32	0.45
4.	T ₄	1.38	0.51
5.	T ₅	1.44	0.58
6.	T ₆	1.26	0.38
7.	T ₇	1.28	0.40
8.	T ₈	1.36	0.49
9.	T ₉	1.40	0.54
10.	T ₁₀	1.22	0.34
	S.E ±	0.018	0.026
	C.D. (at 5%)	0.052	0.077

Stalk

The data in regard to N concentration in stalk given in table 2 showed linear and significant increase in all the treatments over control. Maximum N concentration in stalk was (0.58%) was noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) followed by (0.54%) with T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum (0.34%) at control (T₁₀). Variation in N concentration within 75% RDN, 100% RDN and 125% RDN was recorded non-significant. Integration of S influenced N concentration higher in comparison to FYM and Zn when applied with 75% RDN and 100% RDN treatments.

Phosphorus concentration**Grain**

Data pertaining to P concentration in grain given in table 3 and revealed that P accumulation in grain influenced significantly in all the treatment over control. P concentration in grain varied from 0.25% to 0.37% lowest at control (T₁₀)

and maximum at T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) respectively. Integration of S, Zn and FYM showed slight increase when applied with 100% RDN and 75% RDN treatments. Variation in P concentration within 75% RDN, 100% RDN and 125% RDN was noted narrower and non-significant.

Table 3: Impact of Integrated Nutrient Management on P (%) content in grain and stalk

S. No.	Treatments	P content (%) in grain	P content (%) in stalk
1.	T ₁	0.32	0.128
2.	T ₂	0.30	0.122
3.	T ₃	0.31	0.126
4.	T ₄	0.34	0.134
5.	T ₅	0.37	0.142
6.	T ₆	0.28	0.116
7.	T ₇	0.29	0.120
8.	T ₈	0.33	0.130
9.	T ₉	0.31	0.136
10.	T ₁₀	0.26	0.110
S.E. ±		0.018	0.
C. D. (at 5%)		0.053	0.077

Stalk

P concentration in stalk depicted in table 3 showed linear and non-significant variation within all the treatments. Maximum absorption of P in stalk (0.142%) was noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum (0.110%) at control (T₁₀). Addition of FYM, S and Zn also showed slight increase in P concentration in stalk when applied with 75% RDN and 100% RDN but the increase in P concentration in stalk was noted non-significant. Variation in P accumulation in stalk within 75% RDN, 100% RDN and 125% RDN was also found non-significant.

Potassium concentration in maize

Grain: The data presented in table 4 showed significant influenced of K concentration in grain in all the treatments over control. Maximum K concentration 0.44% was recorded with T₅(100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) followed by 0.42% with T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹)and minimum 0.32%at control (T₁₀). Addition of FYM, sulphur and zinc showed significant influenced in K accumulation at 75% RDN and 100% RDN treatments. Variation in K accumulation in grain within 75% RDN, 100% RDN and 125% RDN was noted non-significant.

Table 4: Impact of Integrated Nutrient Management on K (%) Content in grain and stalk

S. No.	Treatments	K. content in (%) grain	K. content(%) in stalk
1.	T ₁	0.38	1.12
2.	T ₂	0.36	1.09
3.	T ₃	0.37	1.11
4.	T ₄	0.41	1.15
5.	T ₅	0.44	1.19
6.	T ₆	0.34	1.05
7.	T ₇	0.35	1.07
8.	T ₈	0.38	1.13
9.	T ₉	0.42	1.16
10.	T ₁₀	0.32	1.02
S. E. ±		0.021	0.003
C. D. (at 5%)		0.062	0.010

Stalk: Data in respect to K concentration in stalk was given in table 4 showed significant increase in K absorption in all the treatment in comparison to control. Maximum value of K concentration was noted 1.19% in T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 1.02% at control (T₁₀). Variation in P accumulation within 75% RDN, 100% RDN and 125% RDN was noted non-significant. Integration of S, Zn and FYM showed positive influenced in K concentration when applied with 75% RDN and 100% RDN treatments.

that S concentration in all the treatments influenced significantly in over to control. Higher accumulation of S in grain was recorded 0.390% with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 0.20% with T₁₀ (control). Integration of sulphur, Zn and FYM influenced significantly S concentration in grain when applied with 75% RDN and 100% RDN. It was also observed that addition of S showed higher increase in its accumulation when applied with 75% RDN and 100% RDN. Variation in S concentration within 75% RDN, 100% RDN and 125% RDN was also noted significant.

Sulphur concentration in maize

Grain: It is apparent from the data given in table 5 showed

Table 5: Impact of Integrated Nutrient Management on S (%) content in grain and stalk

S. No.	Treatments	S content (%) in grain	S content (%) in stalk
1.	T ₁	0.34	0.27
2.	T ₂	0.32	0.24
3.	T ₃	0.33	0.26
4.	T ₄	0.36	0.30
5.	T ₅	0.39	0.33
6.	T ₆	0.30	0.22
7.	T ₇	0.31	0.23
8.	T ₈	0.35	0.28
9.	T ₉	0.37	0.31
10.	T ₁₀	0.28	0.20
S. E. ±		0.018	0.016
C. D. (at 5%)		0.055	0.047

Stalk

Data in regard to S concentration in stalk given in table 5 showed that accumulation of sulphur in all the treatments influenced significantly over control. Addition of S, Zn and FYM also increase the S concentration when applied with 100% RDN and 75% RDN treatments. Maximum S concentration 0.33% was recorded with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 0.20% at control (T₁₀). Significant variation in S accumulation was also noted within 75% RDN, 100% RDN and 125% RDN treatments.

Zinc concentration in maize

Grain: A critical perusal of the data given in table 6 showed that zinc absorption in grain increase significantly in all the treatments over its control (T₁₀). Highest zinc concentration in grain was noted 31.8 ppm with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and lowest 22.2 ppm with T₁₀ (control). Integration of sulphur, zinc and FYM also showed significant increase when added with 75% RDN and 100% RDN treatments. It was also observed that integration of zinc showed increase in its concentration when applied with 100% RDN and 75% RDN treatments. Variation in zinc concentration within 75% RDN and 100% RDN was found non-significant while variation with 100% RDN and 125% RDN was noted significant.

Stalk

Data presented in table 6 showed significance influenced in Zn concentration in stalk in all the treatments over its

respective control. Maximum zinc concentration 23.8 ppm was noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 14.2 ppm at control (T₁₀). Addition of sulphur, Zn and FYM also influenced zinc accumulation significant when applied with 75% RDN and 100% RDN treatments. Like-wise sulphur addition of zinc also showed higher increase in its concentration in stalk when applied with 75% RDN and 100% RDN treatments. Non-significant accumulation of zinc was noted within 75% RDN, 100% RDN and 125% RDN treatments.

Table 6: Impact of Integrated Nutrient Management on Zn (ppm) content in grain and stalk

S. No.	Treatments	Zn content (ppm) in grain	Zn content (ppm) in stalk
1.	T ₁	27.00	18.80
2.	T ₂	24.80	16.80
3.	T ₃	26.20	18.20
4.	T ₄	29.80	20.80
5.	T ₅	31.8	22.80
6.	T ₆	23.50	15.40
7.	T ₇	24.20	16.20
8.	T ₈	28.60	19.40
9.	T ₉	30.60	21.20
10.	T ₁₀	22.20	14.20
S.E ±		1.509	1.053
C.D. (at 5%)		4.519	3.153

Effect of INM on nutrient uptake**Nitrogen uptake****Grain**

Data in regard to N uptake in grain presented in table 7 revealed that N uptake increased significantly in all the treatment in comparison to its respective control. Variation in N uptake within all the treatment was noted narrower and non significant. Maximum N uptake 50.75 kg ha⁻¹ was recorded with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) followed by 45.63 kg ha⁻¹ with T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 22.87 kg ha⁻¹ at control (T₁₀). Addition of FYM, S and Zn with 100% RDN and 75% RDN also influenced N accumulation. Addition of S with 100% RDN and 75% RDN showed significant accumulation of N uptake while addition of Zn influenced significantly N uptake when applied with 100% RDN. Variation in N uptake within 75% RDN, 100% RDN and 125% RDN was found non-significant.

Table 7: Impact of Integrated Nutrient Management on N uptake in grain and stalk

S. No.	Treatments	N uptake (kg ha ⁻¹) in grain	N uptake (kg ha ⁻¹) in stalk	Total uptake (kg ha ⁻¹)
1.	T ₁	39.12	39.35	78.47
2.	T ₂	34.96	34.11	69.07
3.	T ₃	38.24	38.55	76.79
4.	T ₄	44.84	46.73	91.57
5.	T ₅	50.75	56.83	107.58
6.	T ₆	30.76	28.67	59.43
7.	T ₇	33.91	32.32	66.23
8.	T ₈	40.72	41.96	82.68
9.	T ₉	45.63	49.28	94.91
10.	T ₁₀	22.87	20.40	43.27
S. E. ±		2.14	2.215	4.355
C. D. (at 5%)		6.42	6.632	13.052

Stalk

The data of N uptake in stalk are given in table 7 It is varied

from 20.40 to 56.83 kg ha⁻¹. All the treatment showed significant increase N uptake in stalk over control. Integration

of S and Zn with 100% RDN and 75% RDN showed significant increase in N uptake in stalk. Integration of FYM with 100% RDN and 75% RDN also influenced N uptake in stalk but increase in N uptake in stalk was found non-significant. It was also observed from the data that N uptake in stalk was recorded higher in stalk in comparison to grain but increase N uptake in stalk in comparison to grain was found nominal.

Phosphorus uptake

Grain: It is visualized from the data given in table 8 revealed significant increases in P uptake in grain in all the treatment in comparison to control. P uptake in grain was varied from 4.87 to 13.04 kg ha⁻¹ in all the treatments. Addition of FYM, S and Zn showed significant increase in P uptake in grain when applied with 100% RDN and 75% RDN. Increase in P uptake with 75% RDN and 100% RDN was found significant while increase in P accumulation within 100% RDN and 125% RDN was found non-significant.

Table 8: Impact of Integrated Nutrient Management on P uptake grain and stalk

S. No.	Treatments	P uptake (kg ha ⁻¹) Grain	P uptake (kg ha ⁻¹) stalk	Total uptake (kg ha ⁻¹)
1.	T ₁	9.34	10.94	20.28
2.	T ₂	8.06	9.91	17.97
3.	T ₃	9.00	10.79	19.79
4.	T ₄	11.04	12.28	23.32
5.	T ₅	13.04	13.91	26.95
6.	T ₆	6.81	8.75	15.56
7.	T ₇	7.68	9.69	17.37
8.	T ₈	9.88	11.13	21.01
9.	T ₉	11.40	12.41	23.81
10.	T ₁₀	4.87	6.60	11.47
S. E. ±		0.515	0.600	1.115
C. D. (at 5%)		1.543	1.795	3.338

Stalk

It is appraisal from the data given in table 8 revealed that P uptake in stalk increased significantly in all the treatment over control. Maximum P uptake 13.91 kg ha⁻¹ was noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) followed by 12.41 kg ha⁻¹ with T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 6.60 kg ha⁻¹ at control (T₁₀). Integration of FYM, S and Zn also increased P accumulation when applied with 75% RDN and 100% RDN. P accumulation in stalk within 75%, 100% RDN and 125% RDN was found non-significant. It was also observed that P accumulation in stalk was higher over grain.

Potassium uptake

Grain

It is obvious from the data given in table 9 revealed significant increases in K uptake in all the treatments over control. T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) showed maximum accumulation in 15.51 kg ha⁻¹

followed by T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn) 13.68 kg ha⁻¹ and minimum in 5.99 kg ha⁻¹ in grain was noted with control (T₁₀). Integration of S and Zn showed significant increase in K uptake in grain when applied with 75% RDN and 100% RDN. Addition of FYM also influenced K accumulation in grain but the increase was found non-significant. Variation in K uptake within 75% RDN, 100% RDN and 125% RDN was noted non-significant.

Table 9: Impact of Integrated Nutrient Management on K uptake in grain and stalk

S. No.	Treatments	K uptake kg ha ⁻¹ grain	K uptake (kg ha ⁻¹) stalk	Total uptake (kg ha ⁻¹)
1.	T ₁	11.09	95.82	106.91
2.	T ₂	9.68	88.53	98.21
3.	T ₃	10.74	95.11	105.85
4.	T ₄	13.42	105.39	118.81
5.	T ₅	15.51	116.60	132.11
6.	T ₆	8.27	79.25	87.52
7.	T ₇	9.27	86.49	95.76
8.	T ₈	11.37	96.77	108.14
9.	T ₉	13.68	105.88	119.56
10.	T ₁₀	5.99	61.20	67.19
S. E. ±		0.62	5.191	5.811
C. D. (at 5%)		1.855	15.543	17.398

Stalk

It is apparent from the data given in table 9 showed that K uptake in stalk was significantly increase in all the treatment over control. Highest K uptake 116.60 kg ha⁻¹ was noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) followed by 105.88 kg ha⁻¹ T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 61.20 kg ha⁻¹ at control. Addition of sulphur, Zn and FYM also influenced K accumulation in stalk when applied with 100% RDN and 75% RDN but the increase in K uptake in stalk was found non-significant. Increase in K uptake within 75% RDN, 100% RDN and 125% RDN was also found non-significant. It is also obvious from the data that stalk showed more than 7 times higher accumulation of K in comparison to grain.

Sulphur uptake

Grain

Data in respect to S uptake in grain are given in table 10 showed significant increase in S uptake in grain in all the treatments over control. Addition of sulphur and zinc showed significant increase in S uptake in grain when applied with 75% RDN and 100% RDN. Integration of FYM showed non-significant increase in S accumulation in grain when applied with 75% RDN and 100% RDN. Sulphur accumulation in grain with 75% RDN and 100% RDN was noted non-significant while absorption of S in grain within 100% RDN and 125% RDN was observed significant. Maximum S uptake in grain 13.74 kg ha⁻¹ with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum S uptake 5.24 kg ha⁻¹ with control (T₁₀).

Table 10: Impact of Integrated Nutrient Management on S uptake in grain and stalk

S. No.	Treatments	S uptake (kg ha ⁻¹) grain	S uptake (kg ha ⁻¹) stalk	Total uptake (kg ha ⁻¹)
1.	T ₁	10.92	23.09	33.01
2.	T ₂	8.60	19.49	28.09
3.	T ₃	9.58	22.27	31.85
4.	T ₄	11.69	27.49	39.18

5.	T ₅	13.74	32.33	46.07
6.	T ₆	7.30	16.60	23.90
7.	T ₇	8.21	18.58	26.79
8.	T ₈	10.48	23.97	34.45
9.	T ₉	12.06	28.29	40.35
10.	T ₁₀	5.24	12	17.24
S. E. \pm		0.535	1.357	1.892
C. D. (at 5%)		1.602	4.64	6.242

Stalk

The uptake data of sulphur in stalk given in table 10 showed significant increase in S uptake in all treatment over control. Higher absorption of S in stalk 32.33 kg ha⁻¹ was noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) followed by 28.29 kg ha⁻¹ with T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn) 28.29 kg ha⁻¹ and minimum 12 kg ha⁻¹ at control. Integration of sulphur and zinc with 75% RDN and 100% RDN showed significant increase in S uptake while integration of FYM showed non-significant increase in S uptake when applied with 75% RDN and 100% RDN. Variation in S uptake in stalk within 75% RDN, 100% RDN and 125% RDN was also noted non-significant. It is also obvious from the data that S accumulation in stalk was recorded more than 2 times higher in over grain.

Zinc uptake

Grain

Data in regard to zinc uptake in grain depicted in table 11 revealed that Zn uptake increased significantly in all the treatments over control. Highest Zn uptake 1120.95 gm ha⁻¹ was noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 415.24 gm ha⁻¹ with control (T₁₀). Addition of S and Zn influenced significantly Zn accumulation when applied with 100% RDN and 75% RDN but the increase in Zn uptake in grain was found non-significant. Addition of FYM also influenced Zn accumulation in grain but the increase in Zn uptake in grain was found non-significant. Variation in Zn uptake within 75% RDN 100% RDN and 125% RDN was also noted non-significant.

Table 11: Impact of Integrated Nutrient Management on Zn uptake in grain and stalk

S. No.	Treatments	Zn uptake (gm ha ⁻¹) grain	Zn uptake (gm ha ⁻¹) stalk	Total uptake (gm ha ⁻¹)
1.	T ₁	788.39	1608.33	2396.73
2.	T ₂	661.73	1364.66	2026.39
3.	T ₃	761.10	1559.88	2320.98
4.	T ₄	968.49	1906.31	2874.80
5.	T ₅	1120.94	2234.16	3355.10
6.	T ₆	572.22	1162.38	1734.60
7.	T ₇	641.30	1309.28	1950.58
8.	T ₈	856.56	1661.60	2518.16
9.	T ₉	997.55	1935.10	2932.65
10.	T ₁₀	416.24	852.00	1268.24
S. E. \pm		45.24	91.065	136.30
C. D. (at 5%)		135.399	272.663	408.062

Stalk: It is visualised from the data given in table 11 showed that all the treatment significantly influenced Zn uptake in stalk over control. Highest Zn uptake in stalk was noted 224 gm ha⁻¹ with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum 852 gm ha⁻¹ at control (T₁₀). Addition of S and Zn showed significance increased in Zn uptake when

applied with 100% RDN and 75% RDN while integration of FYM showed non-significant increase in Zn accumulation when added with 100% RDN and 75% RDN. Variation in Zn uptake within 100% RDN and 125% RDN was found significant. It is also obvious from the data that stalk showed more than 2 times higher Zn accumulation over the grain.

Discussion

Impact of INM on nutrient content and uptake

Concentration of nutrient *i.e.* N, P, K, S and Zn were analyzed at harvest in grain and stalk. Based on yield and nutrient concentration their uptake values were also calculated to obtained definite amount of nutrient taken by crop to attain the observed yield level.

Nutrient content

Impact of INM on nutrient content at harvest in grain and stalk presented in table 2 to table 6 that contents of N, P, K, S and Zn in grain and stalk increased significantly in all the treatments over control. Variation in N, P, K, S and Zn content in grain and stalk within all the treatments were noted narrower and non-significant. Addition of S and Zn and FYM with 100% RDN and 75% RDN treatments also influenced N, P, K, S and Zn content in grain and stalk. It was also observed that integration of S and Zn showed higher increase in its concentration and also increase the concentration of other nutrient. The higher concentration of N, P, K, S and Zn to be attributed to higher availability and synergistic effects of these nutrients to each other at all the stage of crop growth. It was also observed that concentration of the nutrient bearing potash was higher in grain than stalk which ascribed to the translocation of N, P, S and Zn from vegetative part of crop plant to grain at the time of maturity. Unlike the higher concentration of N, P, S and Zn in grain content of K reported higher in stalk, this may be due to the fibrous cells which are usually in sclerenchyma cells and they tend to respond to potassium supply. K supply results in relatively high turgidity and high content of cellulose and hemicelluloses associated with high content of K in stalk. Similar results are also reported by other workers Karki *et al.* (2005), Shashidhar *et al.* (2009), Mann *et al.* (2006), Mahala *et al.* (2006) and Vikas *et al.* (2007) [4, 19, 11, 10, 24].

Nutrient uptake

Impact of INM on nutrient uptake at harvest in grain and stalk were computed and presented in table 7 to table 11 it is clearly revealed that all the treatment showed significant increase in N, P, K, S and Zn uptake in grain and stalk over control. Maximum accumulation of N, P, K, S and Zn were noted with T₅ (100% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) followed by T₉ (75% RDN + 25% N FYM + 30 kg S + 5 kg Zn ha⁻¹) and minimum at control (T₁₀). Addition of S, Zn and FYM also influenced N, P, K, S and Zn accumulation in grain and stalk when applied with 100% RDN and 75% RDN.

It was also observed that uptake of N, P, S and Zn was in general higher in stalk, while N, P, S and Zn content was recorded higher in grain. It may be due to the more than 3 times more stalk yield over grain yield. It was also observed that K uptake in stalk was more than 6 times over than grain. It may be due to higher concentration of K in stalk and more than 3 times over stalk yield than grain. These results are in close conformity with the findings of Sujatha *et al.* (2008)^[21], Kumar (2008)^[7], Thavaprakash and Velayudham, (2007)^[22] and Sarwar *et al.* (2012)^[17].

References

- Chesnin L, Yein CH. Turbidiometric method of sulphate sulphur in plant materials. In Proceedings Soil Science Society America. 1951;15:149-151.
- Chandel RS. Electrode melting and plate melting efficiencies of submerged arc welding and gas metal arc welding. Materials science and technology. 1990;6(8):772-777.
- Jackson KA, Uhlmann DR, Hunt JD. On the nature of crystal growth from the melt. Journal of Crystal Growth. 1967;1(1):1-36.
- Karki TB, Ashok Kumar, Gautam RC. Influence of integrated nutrient management on growth, yield, content and uptake of nutrients and soil fertility status in maize (*Zea mays* L.). Indian J Agric. Sci. 2005;75(10):682-685.
- Kumar Prasanna, Halepyati AS, Pujari BT, Desai BK. Effect of integrated nutrient management on productivity, nutrient uptake and economics of maize (*Zea mays* L.) under rainfed condition. Karnataka Journal of Agricultural Sciences. 2007;20:03.
- Kumar V, Singh AP. Long-term effect of green manuring and farmyard manure on yield and soil fertility status in rice-wheat cropping system. Journal of the Indian Society of Soil Science. 2010;58:409-412.
- Kumar A, Singh R, Rao LK, Singh UK. Effect of integrated nitrogen management on growth and yield of maize (*Zea mays* L.) cv. PA-711Madras Agric. J. 2008;95(7-12):467-472.
- Laxminarayana K. Effect of integrated use of inorganic and organic manures on soil properties, yield and nutrient uptake of rice in Ultisols of Mizoram. Journal of the Indian Society of Soil Science. 2006;54:120-123.
- Lindsay WL, Norvell W. Development of a DTPA soil test for zinc, iron, manganese, and copper. Soil science society of America journal. 1978;42(3):421-428.
- Mahala HL, Shaktawat MS, Shivran RK. Direct and residual effect of sources and levels of phosphorus and farmyard manure in maize (*Zea mays* L.) -mustard (*Brassica juncea*) cropping sequence. Indian Journal of Agronomy. 2006;51:10-13.
- Mann KK, Brar BS, Dhillon NS. Influence of long-term use of farmyard manure and inorganic fertilizers on nutrient availability in a Typic Ustochrept. Indian J Agric. Sci. 2006;76(8):477-480.
- Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate (No. 939). US Department of Agriculture, 1954.
- Piper CS. Soil and plant analysis, Hans. Pub. Bombay. Asian Ed, 1966, 368-374.
- Pratt P, Chapman H. Gains and losses of mineral elements in an irrigated soil during a 20-year lysimeter investigation. Hilgardia. 1961;30(16):445-467.
- Ramesh P, Panwar NR, Singh AB, Ramana S. Effect of organic manures on productivity, nutrient uptake and soil fertility of maize (*Zea mays*)- linseed (*Linum usitatissimum*) cropping system. Indian Journal of Agricultural Sciences. 2008;78(4):77-80.
- Roul PK, Sarawgi SK. Effect of integrated nitrogen nutrition techniques on yield, nitrogen content, uptake and use efficiency of rice (*Oryza sativa* L.) Indian Journal of Agronomy. 2005;50(2):129-131.
- Sarwar M, Jilani G, Rafique E, Akhtar EM, Chaudhry AN. Impact of integrated nutrient management on yield and nutrient uptake by maize under rain-fed conditions. Pakistan J Nutri. 2012;11(1):991-998.
- Setia RK, Sharma KN. Dynamics of forms of inorganic phosphorus during wheat growth in a continuous maize-wheat cropping system. Journal of the Indian Society of Soil Science. 2007;55(2):139-146.
- Shashidhar CU, Veeranna HK, Ramesh YM, Somashekarappa PR, Vijay Mahantesh. Effect of different nutrient management practices on yield, economics and nutrient uptake in maize (*Zea mays* L.). Res. on crops. 2009;10(2):27-230.
- Subbaiah BV, Asija GL. A rapid method for estimation of available N in soil. Current Science. 1956;25:259-260.
- Sujatha MG, Lingaraju BS, Palled YB, Ashalatha KV. Importance of integrated nutrient management practices in maize under rainfed condition. Karnataka Journal of Agricultural sciences. 2008;21:334-338.
- Thavaprakash N, Velayudham K. Effect of crop geometry, intercropping and integrated nutrient management practices on cob yield and nutrient uptake of baby corn. Asian Journal of Agricultural Research. 2007;1:10-16.
- Varalakshmi LR, Srinivasamurthy CA, Ann Bhaskar S. Effect of integrated use of organic manures and inorganic fertilizers on organic carbon, available N, P and K in sustaining Productivity of groundnut-finger millet cropping system. Journal of the Indian Society of Soil Science. 2005;53(3):315-318.
- Vikas Abrol, Sharma V, Sharma P, Khar D, Vittal KPR. Direct and residual effect of organic and inorganic sources of nutrients on maize (*Zea mays*)-mustard (*Brassica napus*) cropping sequence under rainfed conditions. Indian Journal of Dryland Agriculture Research and Development. 2007;22(1):82-89.