



ISSN (E): 2277-7695

ISSN (P): 2349-8242

NAAS Rating: 5.23

TPI 2023; 12(5): 849-856

© 2023 TPI

[www.thepharmajournal.com](http://www.thepharmajournal.com)

Received: 28-03-2023

Accepted: 30-04-2023

**Anil Kumar**

Assistant Professor-Cum-Jr.  
Scientist (Soil Science), IRS,  
BAU, Sabour, Bhagalpur, Bihar,  
India

**Diwakar Paswan**

Assistant Professor-Cum-Jr.  
Scientist (Agronomy), JRS,  
BAU, Sabour, Bhagalpur, Bihar,  
India

**SP Vishwkarma**

Assistant Professor-Cum-Jr.  
Scientist (Soil Science), IRS,  
BAU, Sabour, Bhagalpur, Bihar,  
India

**Mrinalini Kumari**

Assistant Professor-Cum-Jr.  
Scientist (Entomology), MBAC,  
BAU, Sabour, Bhagalpur, Bihar,  
India

**Birendra Kumar**

Assistant Professor-Cum-Jr.  
Scientist (Agronomy), BAC,  
BAU, Sabour, Bhagalpur, Bihar,  
India

**Corresponding Author:****Diwakar Paswan**

Assistant Professor-Cum-Jr.  
Scientist (Agronomy), JRS,  
BAU, Sabour, Bhagalpur, Bihar,  
India

## Effect of plant growth promoting rhizobacteria on yield attribute in maize based inter cropping system

**Anil Kumar, Diwakar Paswan, SP Vishwkarma, Mrinalini Kumari and Birendra Kumar**

**Abstract**

48 PGPR isolates were isolated from different rhizotic zones of maize based intercropping system by using different media from twelve different site of Bihar including diara belt. PGPR isolates are coded for *Azospirillum* spp. as AZS<sub>1</sub> to AZS<sub>12</sub>, *Azotobacter* spp. as AZT<sub>1</sub> to AZT<sub>12</sub>, *Pseudomonas* spp. as PSD<sub>1</sub> to PSD<sub>12</sub> and P-solubilizing bacteria spp. as PSB<sub>1</sub> to PSB<sub>12</sub>. These isolates were screened on the basis of seed germination, production of IAA, P-solubilization activity, antifungal activity, and nitrogenase activity for the formulation of microbial consortium. Under pot condition plant height, leaf area index, number of leaves plant<sup>-1</sup>, fresh and dry weight of shoot and root, total biomass production, root volume, percentage root colonization by mycorrhiza in pot soil and microbial population of *Pseudomonas* spp. PSD<sub>6</sub> was maximum in treatment T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) while microbial population of *Azotobacter* spp. AZT<sub>4</sub>, P-solubilization bacteria PSB<sub>4</sub> and MPN of *Azospirillum* spp. AZS<sub>6</sub> were maximum in treatment T<sub>13</sub> (NPK + PSB<sub>4</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>). Keeping in view of experimental findings, PGPRs of diara belt are extremely diversified and perform well in stress condition. They are also competitive in nature and efficient in nitrogen fixing, P-solubilization and producing plant growth hormones. Microbial consortium improves crop growth and increase biomass production. Hence, it may be concluded that PGPRs (PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) form best microbial consortium in all respect than that of others which is very significant not only in growth parameters but also in maintaining soil health for sustainable crop production in Bihar.

**Keywords:** PGPR, rhizobacteria, intercropping, pseudomonas, screening

**Introduction**

Microbial consortium constitutes one of the best possibilities in this aspect, and calls an organized systematic effort for the isolation of different microbial strains including PGPR strains from diverse habitats and developing an effective package of Microbial consortium. A major constraint in exploiting full genetic potential of the crops for achieving higher yield is the supply of adequate nutrients. The application of chemical fertilizer to fulfil the nutrient requirement of crops is advocated since the introduction of green revolution in India.

The rhizobacteria that are beneficial to plant are called plant growth promoting rhizobacteria (PGPR). The term PGPR was first proposed by Kloepper and Schroth (1978) [21] to describe a subset of rhizobacteria which induce increased plant growth after inoculation to seeds. Kloepper (1993) [22] and Cattelan *et al.* (1999) [7] indicated that different strains of PGPR can increase crop yields, control root pathogens, increase resistance against foliar pathogens, promote legume nodulation and enhance seedling emergence. PGPR group of bacteria actively colonize plant roots and increase plant growth and yield (Wu *et al.* 2005) [42]. It may benefit the target plant by causing plant growth promoting and also a source of biological disease control. Growth promoting activity has been reported in strains belonging to several genera, such as *Azotobacter*, *Azospirillum*, *Pseudomonas*, *Acetobacter*, *Burkholderia serratia* and *Bacillus* (Kloepper 1993; Zhang *et al.* 1996, Glick and Bashan 1997, Ramamoorthy *et al.* 2002, Bashan, *et al.* 2004) [22, 44, 14, 45, 5]. Several mechanisms have been postulated to explain the role of PGPR as plant growth, stimulator which can be categorized as direct or indirect promotion. Direct promotion occurs due to ability of targeted strains to produce or change the concentration of phytohormones, like IAA (Mordukhova *et al.* 1991) [27]; Cytokinins (Tien *et al.* 1979) [39], Ethylene (Arshad and Frankenberger 1991, Glick, 1995) [3, 13] and N<sub>2</sub>-fixation by some of the strain (Boddy and Dobereiner, 1995, Mrkovacki and Milic, 2001; Salantur *et al.*, 2006) [6, 28, 46]. Indirect promotion like antagonism against phyto-Pathogenic microorganisms or deleterious bacteria (Kloepper 1993; Glick 1995; Lugtenberg *et al.* 1991) [22, 13, 26] and solubilisation of mineral phosphate and other nutrients (Cattelan *et al.* 1999) [7].

Numerous plant growth promoting rhizobacteria of the genus *Pseudomonas*, *Bacillus* (PSB), *Arthrobacter*, *Azospirillum*, *Klebsiella*, *Azotobacter* and *Enterobacter* have been isolated from the rhizosphere of various crops and have been evaluated for their synergistic effect on plant growth (Klopper *et al.*, 1992) [20]. Egamberdiyeva. D. (2007) [47] studied the rhizosphere and phyllosphere bacteria isolated from wheat and peas and examined for their plant growth promoting properties. Bacterial strains were identified as *Pseudomonas*, *Bacillus* and *Microbacterium* species. After inoculation with effective bacterial strains, the root and shoot growth, and nodulation of peas increased. Gholami *et al.* (2009) [12] also reported that under *In vitro* condition seed treatment with PGPR strains improved seed germination, seedling vigour, seedling emergence and seedling stand over the control in maize.

Maize is grown almost all states of India. Bihar being one of the most important maize growing state ranked fourth in area (6.4 lakh ha) and second in production (17.20 lakh tonnes) in the country in 2006-07. In Bihar maize is grown in all three cropping seasons out of which *Rabi* maize rank first in terms of productivity. Area under winter maize is increasing at a faster rate especially under north Bihar condition. *Rabi* crop is sown in Oct.-Nov; makes little growth till mid-January, leaving enough space for inter cropping during the cropping period. It is planted in rows 60 cm apart. Legumes, Potato, turmeric, tobacco are some successful intercrops that are taken with *Rabi* maize.

## Material and Methods

This chapter deals with the description of the material used and the methods or techniques adopted during the course of investigation.

### 1. Isolation of Plant Growth Promoting Rhizobacteria (PGPR)

Different PGPR isolates were isolated by using selective or non selective media. 10 g soil from the rhizosphere of host plant from selected site were taken and prepared 10 fold dilution series up to  $10^{-6}$  by serial dilution method. 1 ml suspension from different dilutions ( $10^{-2}$  to  $10^{-6}$ ) was poured on plates containing respective media. The suspensions were spread on the plate by using sterilized spreader under aseptic condition.

### 2. Medium for PGPR

Different selective and non selective media were used for isolation of Plant Growth Promoting Rhizobacteria (PGPR). Selective media used were King' B (KB) for *Pseudomonas* spp. Pikovskay' agar (PKV) for PSB, Jensen's medium (JEN) for *Azotobacter* spp., *Pseudomonas fluorescens* agar (PFA) for *Pseudomonas* spp. and N-free malate medium (Nfb) for *Azospirillum* spp. isolates and Non-selective media used for PGPR were nutrient agar (NA), Potato dextrose agar (PDA) and soil extract agar (SEA). Seeds of Maize CV Laxmi were obtained from All India Coordinated Maize Improvement Project (AICMIP), Department of Genetic and Plant Breeding, Tirhut College of Agriculture, Dholi, R.A.U., Pusa, Samastipur, Bihar.

### 3. Studies on survival of PGPR isolates in consortia

A modified succinate broth (MSB) was prepared to grow PGPR isolates, PSD<sub>6</sub>, PSB<sub>4</sub>, AZT<sub>4</sub> and AZS<sub>6</sub> together to

examine their interaction *in vitro* condition.

Modified succinate medium broth (Prasad *et al.*, 1991) [43].

Constituent's	g/L
Sodium glutamate	1.0 g
Sodium succinate	5.0 g
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	0.1 g
MgSO <sub>4</sub> . 7H <sub>2</sub> O	0.1 g
K <sub>2</sub> HPO <sub>4</sub>	0.5 g
Mannitol	10.0 g
NaCl	0.1 g
Yeast Extract	0.5 g
Distilled water	1000.0 ml
pH	6.8

For study of survival of AZT<sub>4</sub> and AZS<sub>6</sub> with PSB<sub>4</sub> and PSD<sub>6</sub> in culture broth fresh inoculum of AZT<sub>4</sub> and AZS<sub>6</sub> were prepared by growing in Jensen's broth and N-free bromothymol blue medium respectively for 96 hrs and PSB<sub>4</sub> and PGPR by growing in MSB was 48 hrs.

50 ml of MSB was taken in 250 ml Erlenmeyer conical flasks, plugged with cotton and sterilized at 103.4 kilo Pascal pressure for 30 min. After cooling the broth medium was inoculated with the 1 ml inoculums of AZT<sub>4</sub>, AZS<sub>6</sub>, PSB<sub>4</sub> and PSD<sub>6</sub>, either alone or in combination according to treatments given below. The inoculated flasks were incubated for  $30 \pm 1$  °C for 6 days. Inoculated flasks were shaken intermittently at  $30 \pm 1$  °C on shaker for ½ hrs 3-4 times daily.

Population of the microorganisms in their inoculums were determined after 3 and 6 days of incubation. For this one ml bacterial suspension from each flask was aseptically transferred in test tubes containing 9 ml sterilized water. Serial dilutions were prepared upto  $10^{-12}$ . The population of AZT<sub>4</sub> was estimated in Jensen's medium, AZS<sub>6</sub> in N-free bromothymol blue medium, PSB<sub>4</sub> in nutrient medium, while PSD<sub>6</sub> in modified succinate agar medium by serial dilution and plate count method.

### 3.1 Germination count

Seed of maize CV laxmi were surface sterilized with 2 per cent sodium hypochlorite for 30 minute followed by washing with 0.01N HCl to avoid its any harmful effect. After proper washing with sterilized distilled water. Seeds were inoculated with different combination of selected PGPR isolates (*Pseudomonas* spp. PSD<sub>6</sub>, *Azotobacter* spp. AZT<sub>4</sub>, *Azospirillum* spp. AZS<sub>6</sub> and *P-solubilizing* bacteria spp. PSB<sub>4</sub>). The inoculated seeds of maize were kept for germination on moist filter paper with the help of sterilized forcep. The inoculated seeds were allowed to germination in incubator at  $28^{\circ} \pm 2$  °C for 7 days according to the following treatment combination in triplicates.

After completion of incubation period the seedlings were carefully taken from the petriplates. The seedlings were then kept on the blotting paper sheet to remove any excess of media and observation regarding shoot and root length measurements were made. In control plate, the seeds treated with sterilized media were taken vigor index was determined with following formula.

Vigor index = (Mean root length + mean shoot length) x germination %.

A pot culture experiment was conducted at department of

Microbiology, F.B.S. and H., R.A.U., Pusa during the Rabi season 2009.

### 3.2 Soil characteristics

Soil was taken from 0-15 cm depth randomly selected spots from Kitchen Garden, R.A.U. Pusa with the help of soil auger. The soil samples were mixed and make composite sample, which was subsequently stirred, dried at room temperature, powdered and finally sieved and kept in polyethylene bag for physico-chemical and microbiological analysis. The soil was found to be low in organic carbon (0.49%), medium in available Nitrogen (250.12 kg/ha), medium in available phosphorus (26.0 kg/ha) and available potash (225.00 kg/ha).

On the basis of morphological, cultural characteristics and screening, four best isolates of PGPR, (*Pseudomonas* spp. PSD<sub>6</sub>, *Azotobacter* spp. AZT<sub>4</sub>, *Azospirillum* spp. AZS<sub>6</sub> and *P-solubilizing* bacteria spp. PSB<sub>4</sub>) were examined alone or in possible combinations with other isolates under pot experiment during Rabi season, 2009.

### 3.3 Pot culture experiment

The experiment was conducted at Department of Microbiology, F.B.S. & H., R.A.U., Pusa in earthen pots under controlled conditions. Soil samples were collected from Kitchen garden of RAU Pusa Campus. The soils was air dried sieved and kept in polyethylene bag. The earthen pot had capacity to 10 kg soil. 6 kg soil was filled up in each earthen pot. All treatments arranged in 48 pots of 3 replicates with 16 pots/replications. The treatments were arranged on completely randomised design (CRD).

### 3.4 Preparation of seed Inoculation

The maize seeds were inoculated with different possible combination of PGPR inoculation (*Pseudomonas* spp. PSD<sub>6</sub> (10<sup>6</sup> cfu ml<sup>-1</sup>) *Azotobacter* spp. AZT<sub>4</sub> (10<sup>7</sup> cfu ml<sup>-1</sup>), *Azospirillum* spp. AZS<sub>6</sub> (10<sup>7</sup> cfu ml<sup>-1</sup>) and *P-solubilizing* bacteria spp. PSB<sub>4</sub> (10<sup>9</sup> cfu ml<sup>-1</sup>). PGPR inoculants were prepared by growing bacteria in respective media. Mixed inoculation of different combination of these organisms were prepared by mixing equal volumes of culture suspension (10 ml) and inoculated with surface sterilized seeds. A 10 per cent solution of Pharmaceutical grade of gum Arabic served as a sticker for inoculants. Seeds were inoculated with gum Arabic as an adhesive and rolled into suspension of bacteria (10<sup>6</sup> to 10<sup>9</sup> cfu ml<sup>-1</sup>) with sterilized glass rod until uniformly coated. Seeds treated with sterile water and gum Arabic served as the uninoculated control. The viable counts of bacteria per seed were found to be (10<sup>6</sup> to 10<sup>8</sup>). Basal doze of fertilizer N @ 200 kg ha<sup>-1</sup>, P<sub>2</sub>O<sub>5</sub> 80 kg ha<sup>-1</sup>, and K<sub>2</sub>O 80 kg ha<sup>-1</sup> in the form of Urea, SSP and MOP were applied Nitrogen was applied in two split dozes ½ at the time of sowing and rest half at 35 days of sowing. Ten inoculated seeds were sown in each pot and thinned to one plant. The plants were irrigated when required. After 60 days of growth from the date of sowing. Plants were carefully uprooted from each pot, washed in acidified detergent solution followed by distilled water and dried in oven at 65 °C.

### Treatment details

T<sub>1</sub> - Uninoculated Control (UIC)

T<sub>2</sub> - NPK

T<sub>3</sub> - PSD<sub>6</sub>

T<sub>4</sub> - AZT<sub>4</sub>

T<sub>5</sub> - AZS<sub>6</sub>

T<sub>6</sub> - PSB<sub>4</sub>

T<sub>7</sub> - PSB<sub>4</sub> + AZT<sub>4</sub>

T<sub>8</sub> - PSB<sub>4</sub> + AZS<sub>6</sub>

T<sub>9</sub> - PSD<sub>6</sub> + PSB<sub>4</sub>

T<sub>10</sub> - AZT<sub>4</sub> + AZS<sub>6</sub>

T<sub>11</sub> - PSD<sub>6</sub> + AZT<sub>4</sub>

T<sub>12</sub> - PSD<sub>6</sub> + AZS<sub>6</sub>

T<sub>13</sub> - PSB<sub>4</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>

T<sub>14</sub> - PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>

T<sub>15</sub> - PSD<sub>6</sub> + AZS<sub>6</sub> + PSB<sub>4</sub>

T<sub>16</sub> - PSD<sub>6</sub> + AZT<sub>4</sub> + PSB<sub>4</sub>

### Result and Discussion

This study was conducted to assess the effect of either single or in different possible combinations of selected PGPR isolates i.e. diazotroph a N<sub>2</sub>-fixers *Azospirillum* spp. AZS<sub>6</sub>, *Azotobacter* spp. (AZT<sub>4</sub>), biocontrol agent *Pseudomonas* spp. (PSD<sub>6</sub>) and Phosphate solubilizing bacteria PSB<sub>4</sub> on maize plants under pot experiment. Total biomass production, nutrients uptake, microbial population dynamics were also assessed by these PGPR consortia on post harvested soil. There is little information available on the possible synergistic effect of these microbes on maize under pot conditions. PGPR isolates selected for each functions were combined together and their effect evaluated in pot experiment.

### Co-inoculation effect of PGPR on maize Plant Physiology Plant height

Plant height of Maize at 45 and 60 DAS increased significantly over uninoculated control (UIC). At 30 DAS PGPR consortia treatment T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZS<sub>6</sub>) gave highest Plant height (11.60 cm) followed by T<sub>15</sub> (NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + PSB<sub>4</sub>) (11.57 cm). The data were non-significant. At 45 day after sowing (DAS) highest plant height was recorded under treatment T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) 35.33 cm which was at par with T<sub>15</sub> and T<sub>16</sub>. The lowest plant height was recorded in UIC (T<sub>1</sub>) 22.23 cm. Mixed inoculation or consortium T<sub>14</sub> significantly superior over single and double inoculants.

At 60 day after sowing (DAS) highest plant height was recorded in T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>) i.e. 65.23 cm which was significantly superior over T<sub>15</sub>, T<sub>16</sub> and T<sub>13</sub> while lowest plant height was recorded in UIC (T<sub>1</sub>) 31.93 cm. Treatment T<sub>14</sub> significantly superior over single and double inoculants. As compared with UIC, T<sub>14</sub> recorded 51.05%, T<sub>15</sub> 49.34% and T<sub>16</sub> 48.14% increased in plant height. This finding was supported by Yaseri and Patwardhan (2007) reported that application of *Azotobacter* and *Azospirillum* strains increased canola yield 21.17% number of branches 11.78%. There are numerous report indicating beneficial effect of *Azospirillum* inoculation in crop yield. Similarly promotion in plant height, plant dry weight and grain yield of various crop plants to inoculation with PGPR were reported by other workers (Khalid *et al.* 2004, Biswas *et al.* 2000a,b) [18, 48].

### Leaf area index

Leaf area index at 45 and 60 DAS increased significantly over uninoculated control (UIC). At 30 DAS leaf area index was non-significant however T<sub>14</sub>, T<sub>15</sub>, T<sub>16</sub> recorded highest value. At 45 DAS highest leaf area index of plant was recorded in

treatment T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>) (0.25) which was at par with treatment T<sub>15</sub> (0.24) and significantly superior over T<sub>16</sub> (0.22) and T<sub>13</sub> (0.22) PGPR consortia. The lowest plant leaf area index was recorded in UIC (T<sub>1</sub>) 0.09. Mixed inoculation or consortium T<sub>14</sub> significantly superior over single and double inoculants. At 60 DAS highest leaf area index of maize plant was recorded in treatment T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) (0.46), which was at par with treatment T<sub>15</sub> (0.43) and significantly superior over T<sub>16</sub> (0.42) and T<sub>13</sub> (0.41) PGPR consortia, while lowest plant leaf area index was recorded in UIC (T<sub>1</sub>) 0.18.

### Number of leaf plant<sup>-1</sup>

Number of leaf plant<sup>-1</sup> in maize crop at 45 and 60 DAS increased significantly over the UIC (T<sub>1</sub>). At 30 DAS number of leaf plant<sup>-1</sup> was non-significant. At 45 DAS highest number of leaf plant<sup>-1</sup> in maize were recorded in consortia consists of triple inoculation treatment T<sub>13</sub>, T<sub>14</sub>, T<sub>15</sub> and T<sub>16</sub> and double inoculation treatment T<sub>8</sub> and T<sub>10</sub> (7.00), which were significantly superior over UIC (T<sub>1</sub>) (5.0) and fertilizer treatment T<sub>2</sub> (5.33).

At 60 DAS highest number of leaf plant<sup>-1</sup> in maize were recorded in consortia consists of triple inoculation treatment T<sub>13</sub>, T<sub>14</sub>, T<sub>15</sub> and T<sub>16</sub> (11.00) which were significantly superior over UIC (T<sub>1</sub>) (7.33) and fertilizer treatment T<sub>2</sub> (7.67). Present studies revealed that maize seed inoculated with all PGPR consortia resulted in an increased plant height, leaf area index and number of leaf plant<sup>-1</sup>. The enhancing effect of seeds inoculation with rhizobacteria applied either double or triple as compared to single inoculant on plant height, leaf area index and number of leaf plant<sup>-1</sup> might be attributed to nitrogen fixing and phosphate solubilizing ability of these micro organisms to produced growth promoting substances. Similar increases in plant height, leaf area, were observed in different crops inoculated with *Pseudomonas*, *Azospirillum* and *Azotobacter* strain (Martinez-Toledo *et al.* 1988; Siddiqui and Shaikat, 2002, Burd *et al.* 2000) [49, 50, 51]. In comparison to *Azotobacter* isolates (AZT<sub>4</sub>), *Azospirillum* isolates (AZS<sub>6</sub>) was found superior, this was because of sandy clay nature of soil, which suited more to *Azospirillum* than *Azotobacter*. Srinivasan (1989) has suggested that *Azospirillum* was more effective in loam to clay soils with high moisture levels, while *Azotobacter* was more effective in semi dry loam. Mixed inoculation of *Azotobacter* and *Azospirillum* were always superior at all N-levels compared to individual application in sugarcane as reported by Navale *et al.* (1985) [52] in extensive trials for three year. Mixed co-inoculation of diazotrophic isolates *Azotobacter* and *Azospirillum* (AZT<sub>4</sub> + AZS<sub>6</sub>) and bio control agent isolates *Pseudomonas* PDS<sub>6</sub> proved to enhance root and foliage growth and to produce significantly plant height, leaf area index and number of leaf plant<sup>-1</sup> than the UIC. Mixed culture of PGPR consortia i.e. AZS<sub>6</sub> + PSD<sub>6</sub> + AZT<sub>4</sub> produced significant increase in the uptake of mineral nutrients over UIC. This increase may be related to the production of hormones by *Azospirillum* isolates (Tie *et al.* 1979) such as auxin, gibberelline and cytokinins. *Pseudomonas* (PSD<sub>6</sub>) PGPR act as biological control of plant pathogens and improve plant growth by inhibiting pathogens through the synthesis of different compounds (Hill *et al.* 1994) [15]. Badaway *et al.*, (2003) [60] reported that mixed co-inoculation of diazotrophic strains; such as *Azotobacter chroococcum* or *Azospirillum lipoferum* with a phosphate solubilizing strains as *B. Polymyxa*, proved to enhance root

and foliage growth, than the uninoculated control. Afzal and Asghari (2008) [53] reported that dual inoculation of PGPR significantly increased plant height, spike length, root and shoot weight of the test wheat crop. The plant growth promoting rhizobacteria might enhanced plant height and productivity by synthesizing phyto-hormones, increasing the availability of nutrients, facilitating the uptake of nutrients (Burd *et al.* 2000) [51].

Inoculation of seedlings of both fir and spruce with *Pseudomonas* isolates increased the plant height, number of leaf, girth of stem and weight of plants significantly over the control has been reported by Zargar *et al.* (2005) [54]. PGPR consortia which produced high amount of IAA also significantly promoted plant height.

### Co-inoculation effect of PGPR on Dry matter yield production of Maize plant at 60 days growth period

#### Fresh weight of shoot and root

Fresh weight of shoot (g plant<sup>-1</sup>) and root (g plant<sup>-1</sup>) at 60 days growth period varied from 30.73 to 75.80 g plant<sup>-1</sup> and 21.67 to 65.10 g plant<sup>-1</sup> respectively, as affected by different treatments. At 60 days growth period maximum fresh weight of shoot and root was recorded in treatment T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) 75.80 g plant<sup>-1</sup> and 65.10 g plant<sup>-1</sup> respectively, which was at par with T<sub>15</sub>, T<sub>16</sub> and T<sub>13</sub>. The minimum fresh weight of shoot and root was recorded in uninoculated control (UIC) 30.73 g plant<sup>-1</sup> and 21.67 g plant<sup>-1</sup> respectively. Treatment T<sub>14</sub> significantly superior over single inoculant treatment T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub> and double inoculants treatments T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub>, T<sub>10</sub>, T<sub>11</sub> and T<sub>12</sub>.

#### Dry weight of shoot and root (g plant<sup>-1</sup>)

Dry weight of shoot (g plant<sup>-1</sup>) and root (g plant<sup>-1</sup>) at 60 days growth period varied from 3.65 to 7.58 g plant<sup>-1</sup> and 2.60 to 7.47 g plant<sup>-1</sup> respectively, as affected different by treatments. Maximum dry weight of shoot and root were recorded in treatment T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) 7.58 g plant<sup>-1</sup> and 7.47 g plant<sup>-1</sup> respectively, which was at par with T<sub>15</sub>, T<sub>16</sub> and T<sub>13</sub>. The minimum dry weight of shoot and root was recorded in UIC 3.65 g plant<sup>-1</sup> and 2.60 g plant<sup>-1</sup> respectively. The seeds inoculated with *Azospirillum* isolate (AZS<sub>6</sub>) or *Azotobacter* (AZT<sub>4</sub>) or *Pseudomonas* isolate (PSD<sub>6</sub>) or either single or in combination resulted higher dry weight of shoot and root (g plant<sup>-1</sup>) with the UIC. Treatment T<sub>14</sub> significantly superior over single inoculant treatment T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub> and double inoculants, treatment T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub>, T<sub>10</sub>, T<sub>11</sub> and T<sub>12</sub>. Significant increase in shoot and root dry weight were obtained by inoculation with any single isolates, mixed inoculants with two or three isolates as compared to UIC. Mixed inoculants of PGPR consortium consisted of (PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) was superior in plant dry matter (yield) performance. Hussain and Vancura, 1970 [55] reported that inoculation of maize seeds with strains *Pseudomonas fluorescence* and *Chromobacterium violaceum* significantly increased the dry matter yield of plants. Diazotrophs (AZT<sub>4</sub> and AZS<sub>6</sub>) being an efficient N<sub>2</sub>-fixer fixed atmospheric nitrogen to the plant which helps in enhancing photosynthetic efficiency and therefore, there was more accumulation of dry matter. Similar results of increment in dry matter yield of maize due to inoculation of diazotrophs were reported (Rai and Hunt, 1993) [33, 34].

Rhizobacteria (PSD<sub>6</sub>) influence the plant dry matter which probably due to production of plant growth hormones and colonization of beneficial microbes. (Sindhu *et al.*, 1999,

Singh and Pareek 2003) [36, 47].

### Total biomass production

A perusal of data presented in Fig.1 indicated a significant increase in total Biomass production at 60 days growth period which varied from 6.25 to 15.05 g plant<sup>-1</sup> as affected by different treatments. The maximum total biomass production was recorded in the treatment (T<sub>14</sub>) ((NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) 15.05 g plant<sup>-1</sup>, while lowest in the UIC (T<sub>1</sub>) 6.25 g plant<sup>-1</sup>. Treatment T<sub>14</sub> was at par with T<sub>15</sub>, T<sub>16</sub> and T<sub>13</sub> and significantly superior over single inoculant treatment T<sub>3</sub>, T<sub>4</sub>, T<sub>5</sub> and T<sub>6</sub> and double inoculants treatment T<sub>7</sub>, T<sub>8</sub>, T<sub>9</sub>, T<sub>10</sub>, T<sub>11</sub> and T<sub>12</sub>.

The highest biomass production in PGPR consortium T<sub>14</sub> ((NPK + PSD<sub>6</sub> + AZS<sub>6</sub> + AZT<sub>4</sub>) may be due to higher leaf area index and better root and shoot and nutrients from the deeper zone of the soil combined with the better nitrogen fixation capacity of *Azospirillum* (AZS<sub>6</sub>) and *Azotobacter* (AZT<sub>4</sub>) isolates and root colonizer by *Pseudomonas* isolates (PDS<sub>6</sub>). Similar result was reported by Kennedy and Tehan (1992) [61].

The plant growth promoting rhizobacteria play an active role in soil through their natural ability to provide important but scarce nutrients to the plants. The inoculation effect of PGPR including N<sub>2</sub>-fixer AZS<sub>6</sub> and AZT<sub>4</sub> and Root colonizer *Pseudomonas* isolate (PSD<sub>6</sub>) in processed calcareous soils

produced a significant increase in the uptake of mineral nutrients over uninoculated control (UIC). This increase may be related to the production of hormones such as auxin (Mordukhova *et al.* 1991) [27] gibberellins (Mahmaud *et al.* 1984) and cytokinins (Tien *et al.* 1979) [39]. *Azospirillum* and *Pseudomonas* which lives on or in the roots may be responsible for the production of hormones affecting the plant growth by stimulating root growth proliferation of maize. Increased root surface might have resulted in enhanced mineral uptake (Okon 1985) [29]. These observations support the findings of Lin *et al.* 1983) [56] that *A. brasilense* inoculation can improve N uptake of plants and improve the availability and efficiency of applied mineral nutrients. Use of diazotrophs as microbial inoculants has resulted in 20, 15 and 60-80% increase in yield of paddy, wheat and legume respectively and saving of 5-100% chemical fertilizers (Haffeez *et al.* 2002) [57]. Increased in plant height and root and shoot biomass of wheat was reported earlier from twelve different isolates of PGPR. (De Freitas and Germida 1990, 1992) [58, 59]. Several possible mechanism of plant growth promotion have been reported. Among these, the production of biologically active metabolites, particularly the plant growth regulators by rhizosphere microbiota affect plant growth directly after being taken up by the plant or indirectly by modifying the rhizosphere environment (Arshad and Frankenberger, 1998) [3].




Fig 1: Combined effect of plant growth promoting rhizobacteria on Dry matter production of maize crop at 60 days growth period.

### Summary and conclusions

The present investigation was under taken to isolate plant growth promoting rhizobacteria (PGPR) from twelve different diversified area of Bihar including, diara belt (Koshi and Ganga), where microbial diversity was maximum under maize based inter cropping system. Efforts were also made to screen the PGPRs for their plant growth promoting attributes like seed germination, production of I.A.A., P-solubilizing

activity, antifungal activity and nitrogenase activity. The insertion of isolates for the formulation of microbial consortium and its effect on plant physiological parameters, total biomass production, nutrients uptake, and microbial population on post harvest soil were under taken. Selected four best isolates of PGPR *Pseudomonas* (PSD<sub>6</sub>), *P-solubilizer* (PSB<sub>4</sub>), *Azospirillum* (AZS<sub>6</sub>) and *Azotobacter* (AZT<sub>4</sub>) were examined alone or in the possible combinations

with other isolates. The pot experiment was conducted at the Department of Microbiology, F.B.S. & H., R.A.U., Pusa in completely randomised design (CRD) with three replications during rabi 2008-09.

A brief summary of the result on each aspect has been presented given here in following paragraphs.

1. Under pot culture condition plant height of maize at 45 and 60 day after sowing (DAS) was significantly increased by over uninoculated control (UIC). At 45 and 60 DAS, maximum plant height was recorded in PGPR consortia T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>) (35.33 cm and 65.23 cm), followed by T<sub>15</sub> (34.33 and 63.03 cm) and minimum in UIC (T<sub>1</sub>) (22.23 cm and 31.93 cm), respectively.
2. Leaf area index of maize plant at 45 and 60 day after sowing (DAS) was significantly increased over uninoculated control (UIC). At 45 and 60 DAS, maximum leaf area index was recorded in PGPR consortia T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>) (0.25 and 0.46), followed by T<sub>15</sub> (0.24 and 0.43) and minimum (0.09 and 0.18) in UIC (T<sub>1</sub>), respectively.
3. Average number of leaf plant<sup>-1</sup> at 45 and 60 day after sowing (DAS) was significantly increased over uninoculated control (UIC). At 45 and 60 DAS, maximum average number of leaf plant<sup>-1</sup> was recorded in T<sub>14</sub>, T<sub>15</sub>, T<sub>16</sub> and T<sub>13</sub> and minimum in UIC (T<sub>1</sub>) respectively.
4. Fresh weight (g plant<sup>-1</sup>) of shoot and root after 60 days growth period was significantly increased over UIC. Maximum Fresh Weight of shoot and root was recorded in PGPR consortia T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>) (75.80 g plant<sup>-1</sup> and 65.10 g plant<sup>-1</sup>), followed by T<sub>15</sub> (74.33 g plant<sup>-1</sup> and 64.57 g plant<sup>-1</sup>), T<sub>16</sub> (71.67 g plant<sup>-1</sup> and 63.27 g plant<sup>-1</sup>) and T<sub>13</sub> (71.27 g plant<sup>-1</sup> and 62.47 g plant<sup>-1</sup>) and minimum in UIC (T<sub>1</sub>) (30.73 g plant<sup>-1</sup> and 21.67 g plant<sup>-1</sup>) respectively.
5. Dry weight (g plant<sup>-1</sup>) of shoot and root after 60 days growth period was significantly increased over UIC. Maximum dry weight of shoot and root was recorded in PGPR consortia T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>) (7.58 g plant<sup>-1</sup> and 7.47 g plant<sup>-1</sup>), followed by T<sub>15</sub> (7.43 g plant<sup>-1</sup> and 7.52 g plant<sup>-1</sup>), T<sub>16</sub> (7.41 g plant<sup>-1</sup> and 7.05 g plant<sup>-1</sup>) and T<sub>13</sub> (7.34 g plant<sup>-1</sup> and 7.00 g plant<sup>-1</sup>) and minimum in UIC (T<sub>1</sub>) (3.65 g plant<sup>-1</sup> and 2.60 g plant<sup>-1</sup>), respectively.
6. Total biomass production after 60 day growth period was significantly increased over UIC, which was maximum in PGPR consortia T<sub>14</sub> (NPK + PSD<sub>6</sub> + AZT<sub>4</sub> + AZS<sub>6</sub>) (15.05 g plant<sup>-1</sup>), followed by T<sub>15</sub> (14.95 g plant<sup>-1</sup>), T<sub>16</sub> (14.46 g plant<sup>-1</sup>) and T<sub>13</sub> (14.34 g plant<sup>-1</sup>) and minimum in UIC (T<sub>1</sub>) (6.25 g plant<sup>-1</sup>). In the light of summarised result following conclusion emerged.

Under this investigation it was confirmed that maize based cropping system in diara belt was very specific in respect of microbial diversity. This might be due to intercropping with various C<sub>3</sub> species including legumes, C<sub>4</sub> nature and photo insensitiveness of maize crops low fertility status of soil. It was also found that microbial isolates like *Pseudomonas* spp. (PSD<sub>6</sub>), *Azotobacter* spp. (AZT<sub>4</sub>), *Azospirillum* spp. (AZS<sub>6</sub>) and P-solubilizing bacteria spp. (PSB<sub>4</sub>) of these areas are very efficient in biomass production by synthesizing growth hormones in maize based cropping system. On the basis of

screening test, with regard to seed germination test, production of IAA, P-solubilization test, antifungal activity and nitrogenase activity, four most efficient native PGPR isolates of *Pseudomonas* spp. (PSD<sub>6</sub>), *Azotobacter* spp. (AZT<sub>4</sub>), *Azospirillum* spp. (AZS<sub>6</sub>) and P-solubilizing bacteria spp. (PSB<sub>4</sub>) were selected, out of 48 isolates, which were collected from different sites of Bihar. In pot culture experiment maize seeds were inoculated with selected isolates of PGPR alone and different possible consortium and observed enhancement in maize plant physiological attributes, total biomass production and nutrient uptake which were significantly superior over the uninoculated control. The present studies revealed that microbial consortium (*Pseudomonas* PSD<sub>6</sub> + *Azotobacter* AZT<sub>4</sub> + *Azospirillum* AZS<sub>6</sub>) was the best in all respect than that of others and very significant not only in growth parameters but also in maintaining soil health for sustainable crop production. From these result we conclude that the native isolates of PGPR activity isolates from Bihar soil can play an important role in helping plant to establish and growing calcareous condition. It is advisable that isolates may be used in field trial to ascertain weather this combination of these isolates enhancement growth and yield of maize would be feasible.

## References

1. Ahmad F, Ahmad I, Khan MS. Indole Acetic Acid production by the Indigenous Isolates of *Azotobacter* and Fluorescent *Pseudomonas* in the presence and absence of tryptophan. *Truk J Biol.* 2005;29:29-34.
2. Anu Kundu BS. Development and characterization of lacZ marked strains of phosphate solubilizing bacteria. *Indian J. Microbiol.* 2005;45(4):261-263.
3. Arshad M, Frankenberger Jr WT. Influence of ethylene produced by soil microorganisms on etiolated pea seedlings. *Appl. Environ. Microbiol.* 1991;54(11):2728-2732.
4. Aslantas R, Cakmakci R, Sahin F. Effect of plant growth promoting rhizobacteria on young apple trees growth and fruit yield under orchard conditions. *Sci. Hortic.* 2007;111(4):371-377.
5. Bashan Y, Holguin G, De-Bashan LE. *Azospirillum* plant relationships : Physiological, molecular, agricultural and environmental advances (1997-2003) *Con. J Microbiol.* 2004;50(8):521-577.
6. Boddey RM, Dobereiner J. Nitrogen fixation associated with grasses and cereals. Recent progress and perspectives for the future. *Fert. Res.* 1995;42:241-250.
7. Cattelen AJ, Hartel PG, Fuhrmann JJ. Screening for plant growth promoting rhizobacteria to promote early soyabean growth. *Soil Sci. Soc. Am. J.* 1999;63(6):1670-1680.
8. Dobereiner J, Day JM. Associative Symbiosis in tropical grasses: Characterization of micro-organisms and dinitrogen fixing. *Proceedings of First International Symposium on Nitrogen Fixation.* (Eds.) Newtas, W.E. and Nyman, G.J. University of Washington Press, Pullman. 1976;2:518-38.
9. Dunne C, Moenne-Loccoz Y, Me Carthy J, Higgins P, Pawell J, Dowling DN, *et al.* Combining proteolytic and phloroglucinol producing bacteria for improved biocontrol of phythium-mediated damping off of sugar beet. *Plant Pathol.* 1998;47(3):299-307.
10. Forbes BADF, Sahn, Weissfeld AS. Bailey and Scott's

- Diagnositc Microbiolgy. 1998; 11<sup>th</sup> ed., Mosby In c., St, Louis, Missouri, USA.
11. Gaur AC, Pareek KP. Organic acids in soil during degradation of organic residues: Proc. India National Acad. Sci. 1972;40:68.
  12. Gholami A. Shahsavani S, Nezarat S. The effect of plant growth promoting rhizobacteria (PGPR) on germination, seedling growth and yield of maize. International J. Biological and Life Science. 2009;1(1):35-40.
  13. Glick BR. The enhancement of Plant growth by free living bacteria. Con. J. Microbiol. 1995;32:145-148.
  14. Glick BR, Bashan Y. Genetic manipulation of plant growth promotion bacteria to enhance biocontrol of phytopathogens. Biotechnology Advance. 1997;15(2):353-378.  
doi: 10.1016/50734-9750(97)00004-9
  15. Hill DS, Stein JI, Torkewitz NR, Morse AM, Howell CR, Pachalotka JP, *et al.* Cloning of genes involved in the synthesis of Pyrrolinitrin from *Pseudomonas fluorescens* and role of pyrrolnitrin synthesis in biological control of plant disease. Appl. Environ. Microbiol. 1994;60(1):78-85.
  16. Jackson ML. Soil chemical analysis. Prentice Hall (India). Pvt. Ltd., New Delhi; c1993.
  17. Kavimandan SK, Gaur AC. Effect of seed inoculation with *Pseudomonas* spp. on phosphate uptake and yield of maize. Curr. Sci. 1971;40(16):439-440.
  18. Khalid A, Arshad M, Zahiz ZA. Screening plant growth promoting rhizobacteria for improving growth and yield of wheat. Journal of Applied Microbiology. 2004;96(3):473-480.
  19. King EO, Ward MK, Raney DE. Two simple media for the demonstration of Pycocyanin and fluorescein. J. Lab Clin. Med. 1994;44(2):301-307.
  20. Kloepper JM, Kabana RR, Macinroy JA, Young RW. Plant Soil. 1992;139:75.
  21. Kloepper JW, Schroth MN. Plant growth promoting rhizobacteria on radishes. Proc. IV Int. Conf. Plant Pathogen. Bacter. 1978;2:879-882 (c.f. Kumar, S., 2004 M.Sc. Thesis RAU, Pusa).
  22. Kloepper JW. Plant growth promoting rhizobacteria as biocontrol agents. In Blaine Meeting Ir. Ed.; Soil Microbiol. Ecology: Application on agricultural and Environment. Marcell Dekker Inc., New Yark, 1993, 255-274.
  23. Kumar S, Yadav K. Microbial solubilization of inorganic phosphates sources by P-solubilizing micro organisms isolated from calcareous soils. RAU J. Res. 2005;15(1-2):114-117.
  24. Leinhos V, Vasek O. Biosynthesis of auxins by Phosphate solubilizing rhizobacteria from wheat (*Triticum aestivum*) and rye (*Secale cereal*). Microbiol Research. 1994;149(1):31-35.
  25. Li RP, Mac Rac IC. In Proc IX th Australian Nitrogen Fixation Conf. 9:9, c.f. Indian J. Microbiol. 1991;38:196.
  26. Lutenberg BJJ, De Wegez LA, Bennett JW. Microbial stimulation of plant growth and protection from disease. Curr. Opinion Biotechnol. 1991;2(3):457-464.
  27. Mordukhova EH, Skvortsova NP, Kochetkov VV, Dubeikovaskii AN, Boronin AM. Synthesis of the phytohormone indole-3 acetic acid by rhzosphere bacteria of the genus *Pseudomonas*. Mikra *biologiya*. 1991;60:494-500.
  28. Mrekovacki N, Millic V. Use of *Azotobacter chroococcum* as potentially useful in agricultural application. Ann. Microbiol. 2001;51(2):145-158.
  29. Okon Y, Albrecht SL, Burris RH. Methods for growing spirillum lipoferum and for counting it in pure culture and association with plants. Appl. Environ. Microbiol. 1977;33(1):85-87.
  30. Patten CL, Glick BR. Roe of *Pseudomonas putida* indoleacetic acid in development of the host plant root system. Appl. Environ. Microbiol. 2000;68(8):3795-3801.
  31. Picard C, Bosco M. Maize heterosis affects the structure and dynamics of indigenous rhizospheric auxins producing *Pseudomonas* populations. FEMS-Microbiology Ecology. 2005;53(3):349-357.
  32. Prikryl Z, Vanchura V, Wurst M. Auxin formation by rhizosphere bacteria as a factor of root growth. Biol. Plantarum. 1985;27:159-163.
  33. Rai R, Hunt PG. Inoculation of Maize varieties with salt tolerant mutants of *Azospirillum brasilense* and VAM fungi in saline Calcareous soil. Microbial Release. 1993;1:243-251.
  34. Rai R, Hunt PG. Inoculation of Maize varieties with salt tolerant mutants of *Azospirillum brasilense* and VAM fungi in saline Calcareous soil. Microbial Release. 1993;1:243-251.
  35. Ramazan C, Mesude FD, Ummugulsum E. The effect of plant Growth Promoting Rhizobacteria on Barley seedling growth, Nutrient uptake, some soil properties, and bacterial counts. Turk J Agric. 2007;81:189-199.
  36. Sindhu SS, Dadarwal KR, Dahiya BS. Hydrogen evaluation and relative efficiency in chickpea. Effect of Rhizobium strain, Host cultivar and temperature. Indian J Exp. Biol. 1986;24:416-420
  37. Singh RD, Vairanvan K, Ramasamy M. Integrated phosphorus management in green gram. Madras Agric. J. 2002;89:149-150
  38. Sundara Rao WVB, Sinha MK. Phosphate dissolving organisms in the soil and rhizosphere. Indian Journal of Agricultural Science. 1963;33:272-78.
  39. Tien T, Gaskins MH, Hubbell DH. Plant growth substances produced by *Azospirillum brasilense* and their effect on the growth of pearl millet (*Pennisetum americanum* L.) Application and Environmental Microbiology. 1979;37(5):1016-1024.
  40. Vasundhara C, Muracedhara KG, Jacob VB, Sethuraj MR, Kothandoraman R. Nitrogen Fixation by *Azotobacter Chroococcum* under cadmium stress. Indian J. Microbiology. 2002;42(1):15-17.
  41. Voiscard C, Keel C, Haas D, Defago G. Cyanide production by *Pseudomonas fluorescens* helps suppress black root rot of tobacco under genotobiotic conditions. EMBO J. 1989;8(2):351-358.
  42. Wu SC, Cao ZH, Li ZG, Cheung KC, Wong MH. Effects of biofertilizer containing N-fixer, P and K solubilizers and AM fungi on Maize growth: A greenhouse trial. Geoderma. 2005;125(1-2):155-166.
  43. Yadav K, Prasad V, Mandal K. Effect of *Azospirillum lipoferum* on the yield of maize (*Zea mays* L.) genotypes at different levels of nitrogen. RAU J. Res. 1991;9(1&2):44-46.
  44. Zhang T, Ramakrishnan R, Livny M. BIRCH: an efficient data clustering method for very large databases.

- ACM sigmod record. 1996 Jun 1;25(2):103-14.
45. Ramamoorthy V, Raguchander T, Samiyappan R. Induction of defense-related proteins in tomato roots treated with *Pseudomonas fluorescens* Pf1 and *Fusarium oxysporum* f. sp. lycopersici. Plant and soil. 2002 Feb;239(1):55-68.
  46. Salantur A, Ozturk A, Akten S. Growth and yield response of spring wheat (*Triticum aestivum* L.) to inoculation with rhizobacteria. Plant Soil and Environment. 2006 Jan 1;52(3):111.
  47. Egamberdiyeva D. The effect of plant growth promoting bacteria on growth and nutrient uptake of maize in two different soils. applied soil ecology. 2007 Jun 1;36(2-3):184-9.
  48. Biswas JC, Ladha JK, Dazzo FB, Yanni YG, Rolfe BG. Rhizobial inoculation influences seedling vigor and yield of rice. Agronomy Journal. 2000 Sep;92(5):880-6.
  49. Martinez-Toledo MV, De La Rubia T, Moreno J, Gonzalez-Lopez J. Root exudates of Zea mays and production of auxins, gibberellins and cytokinins by *Azotobacter chroococcum*. Plant and soil. 1988 Aug;110:149-52.
  50. Siddiqui IA, Shaukat SS. Mixtures of plant disease suppressive bacteria enhance biological control of multiple tomato pathogens. Biology and Fertility of soils. 2002 Oct;36:260-8.
  51. Burd TD, Pering TA, Stratakos AJ, Brodersen RW. A dynamic voltage scaled microprocessor system. IEEE Journal of solid-state circuits. 2000 Nov;35(11):1571-80.
  52. Navale PA, Mungse HB, Harinarayana G. Combining ability of maintainers in pearl millet. Indian Journal of Agricultural Sciences (India); c1985.
  53. Afzal A, Bano A. Rhizobium and phosphate solubilizing bacteria improve the yield and phosphorus uptake in wheat (*Triticum aestivum*). Int J Agric Biol. 2008 Jan 1;10(1):85-8.
  54. Zargar AH, Singh B, Laway BA, Masoodi SR, Wani AI, Bashir MI. Epidemiologic aspects of postpartum pituitary hypofunction (Sheehan's syndrome). Fertility and sterility. 2005 Aug 1;84(2):523-8.
  55. Hussain A, Vančura V. Formation of biologically active substances by rhizosphere bacteria and their effect on plant growth. Folia microbiologica. 1970 Nov;15:468-78.
  56. Lin YL, Farley RD, Orville HD. Bulk parameterization of the snow field in a cloud model. Journal of Applied Meteorology and climatology. 1983 Jun;22(6):1065-92.
  57. Hafeez K, Zhang Y, Malak N. Core competence for sustainable competitive advantage: A structured methodology for identifying core competence. IEEE transactions on engineering management. 2002 Aug 7;49(1):28-35.
  58. De Freitas JR, Germida JJ. Plant growth promoting rhizobacteria for winter wheat. Canadian journal of microbiology. 1990 Apr 1;36(4):265-72.
  59. De Freitas JR, Germida JJ. Growth promotion of winter wheat by fluorescent pseudomonads under field conditions. Soil Biology and Biochemistry. 1992 Nov 1;24(11):1137-46.
  60. Draz EI, Emara AM, Saad KM, Badaway A. Genotoxicity of some commonly used antidepressants (fluoxetine, sertraline and clomipramine). Mansoura Journal of forensic medicine and clinical toxicology. 2009 Jul 1;17(2):63-78.
  61. Kennedy IR, Tehan YT. Biological nitrogen fixation in non-leguminous field crop, recent advances. Plant Soil. 1992;141:93-118.