



ISSN (E): 2277-7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2023; 12(5): 915-917
© 2023 TPI
www.thepharmajournal.com
Received: 22-02-2023
Accepted: 25-03-2023

Rajeev Singh
Senior Scientist and Head,
Krishi Vigyan Kendra, Manpur,
Gaya, Bihar, India

Nityanand
Senior Scientist and Head Krishi
Vigyan Kendra, Aurangabad,
Bihar, India

Ravi Ranjan Kumar
Subject Matter Specialist
(Ag. Engineering) Krishi Vigyan
Kendra, Aurangabad, Bihar,
India

Tej Pratap
Research Associate, Krishi
Vigyan Kendra, Manpur, Gaya,
Bihar, India

Mohd. Zakir Hussain
Subject Matter Specialist
(Agromet), Krishi Vigyan
Kendra, Manpur, Gaya, Bihar,
India

Bal Manohar
Ph.D Scholar, Department of
Agronomy, Bihar Agricultural
University, Sabour, Bhagalpur,
Bihar, India

Corresponding Author:
Ravi Ranjan Kumar
Subject Matter Specialist
(Ag. Engineering) Krishi Vigyan
Kendra, Aurangabad, Bihar,
India

Effect of sowing time of wheat (*Triticum aestivum* L.) under timely sown condition to avoid terminal heat stress in South Bihar condition

Rajeev Singh, Nityanand, Ravi Ranjan Kumar, Tej Pratap, Mohd. Zakir Hussain and Bal Manohar

Abstract

A field experiment was conducted at Krishi Vigyan Kendra, Aurangabad and in farmers' field during rabi seasons of 2013-14 and 2014-15, to evaluate the suitable time of Wheat (*Triticum aestivum* L.) sowing under timely sown condition to avoid terminal heat in South Bihar. Experiment was laid out in a completely randomized block design with 4 date of sowing. 5th November, 15th November 25th November, 5th December in a total of five replications during rabi 2013-14 and 2014-15. Sowing of wheat at 5th November produce significantly maximum grain yield (44.64 q/ha) being at par with 15th November both were significantly more over 25th November and 5th December. Sowing of wheat at 5th November and 15th November produces 6.13%, 3.70% more grain yield over 25th November and 33.65% and 30.59% more grain yield over 5th December, respectively. Straw yield (52.90 q/ha) recorded maximum with 5th November being at par with 15th November both were significantly more over 25th November and 5th December. Sowing of wheat at 5th November and 15th November produces 8.49%, 7.51% more straw yield over 25th November and 15.96% and 14.91% more straw yield over 5th December. The B-C ratio also recorded significantly higher with 5th November and statically at par with 15th November over 25th November and 5th December. The higher returns by 6.13 and 3.71% & by 33.65% & 30.60%, respectively were recorded when wheat sown at 5th November and 15th November than 25th November and 5th December respectively.

Keywords: Terminal heat stress, wheat, time of sowing

Introduction

Wheat is one of the most staple foods of the humanity (Meena *et al.*, 2013) [5]. Its area and productivity is increasing rapidly adopting across the globe, due to its wider adaptability sustainability under diverse agro climatic conditions (Kumar *et al.*, 2014) [4]. However, considerable portion of the wheat grown in South Asia is considered to be affected by heat stress, of which the majority is present in India (Joshi *et al.*, 2007a) [2]. In India terminal heat stress is a major reason of yield decline in wheat due to delayed planting (Joshi *et al.*, 2007a) [2]. Selection of suitable crop varieties according to the agroclimatic conditions may play crucial role in realizing the optimum production of any crop commodity (Singh *et al.*, 2008). The most heat-stressed locations of South Asia are the Eastern Gangetic Plains (EGP), central and peninsular India, whereas heat stress is considered moderate in north western parts of the Indo-Gangetic Plains (IGP) (Joshi *et al.*, 2007b) [3]. Late planted wheat suffers drastic yield losses which may exceed to 40-50%. Global climate models predict an increase in mean ambient temperature between 1.8 and 5.8 °C by end of this century (IPCC, 2007) [1]. Grain yield was negatively related to the thermal time accumulated above the base temperature of 31 °C (Mian *et al.*, 2007) [6]. High temperature above 32 °C has been reported reducing grain yield and grain weight (Wardlaw *et al.*, 2002) [9]. Shrivelled small grains are produced and different yield associated traits such as tillering, grain weight and grains numbers/spike are reduced. Using this factor (3-4 % loss per 1 °C above 15-20 °C), it can be calculated that most commercially sown wheat cultivars in India would lose approximately 50 % of their yield potential when exposed to 32-38 °C temperature at the crucial grain formation stage. The experiment was conducted at the at Krishi Vigyan Kendra, Aurangabad and farmers of Aurangabad district during the years rabi 2013-14 and 2014-15. By the late sowing the varieties was given high temperature stress during grain filling stage in comparison to timely sown condition.

Materials and Method

The field experiment was conducted at Krishi Vigyan Kendra and farmers' field in Aurangabad district of Bihar during the two consecutive rabi seasons of 2013-14 and 2014-15. The experimental site is situated in South Bihar at 24^o.50' N, 84^o.70' E, and at 332' above mean sea level. The maximum temperature remained above 35.60 °C and 35.97 °C during 2013-14 and 2014-15, respectively. The total rainfall received during crop period was 10.77 and 13.25 mm during 2013-14 and 2014-15, respectively (Fig. 1). The soil was clay-loam having normal soil reaction (pH 7.5), low in organic carbon (0.51%) and available nitrogen (205.7 kg/ha), and medium in available phosphorus (19.3 kg/ha) and available potassium (198.5 kg/ha). The experiment was laid out in completely randomized block design with 5 replications comprising of 4 date of sowing i.e. 5th November, 15th November 25th November, 5th December. In experimental plots, wheat was established by with zero-till drill (ZTD). The wheat variety HD-2733 was tested in different dates of sowing. The fields were leveled with leveler to allow drill to place seeds at

a uniform distance and proper depth in all the replications. The experimental plots meant for zero-till drill (ZTD) sowing were subjected to two ploughing followed by harrowing and planking before sowing with zero-till drill (ZTD) by planking on four date of sowing. Experimental field was fertilized at the rate of 120:60:40 kg NPK/ha. Nitrogen was applied in three splits (1/2 dose of N at basal rest 1/2 dose each equal at 1st irrigation and 2nd irrigation), while the entire P₂O₅ and K₂O were applied as basal application. Pendimethalin was sprayed within 1 days after sowing, by knap-sack sprayer using 800 litres/ha water in all treatment plots in all replications. Post-emergence herbicides, metsulfuron @ 33 g/ha, was applied with knap-sack sprayer fitted with flat-fan nozzle using 500 litres/ha of water at 30 days after sowing (DAS) in all treatment plots in each replications. The data on plant height, number of tillers, crop biomass and number of grains/spike were recorded. The crop was harvested manually in the second week of April. On the basis of existing price of the inputs and outputs, variable cost of cultivation and gross returns were calculated.




Fig 1: Maximum and Minimum Temperature during crop period 2013-14 and 2014-15

Results and Discussion

Number of effective tillers/m², spike length, grains/spike and test-weight were significantly influenced by different date of sowing. Number of effective tillers/m² at harvest stage recorded maximum with wheat sown at 5th November being at par with 15th November both were significantly higher over 25th November and 5th December. Spike length was recorded significantly higher with wheat sown at 5th November over other treatment. Number of grain significantly influenced by date of sowing maximum number of grain/spike was recorded with when wheat was sown at 5th November over 15th November, 25th November and 5th December. 1000 grain weight was also significantly influenced by date of sowing. Maximum 1000 grain weight recorded with wheat sown at 5th November being at par with 15th November they were significantly higher over 25th November and 5th December (Table 1).

Sowing of wheat at 5th November produce significantly maximum grain yield (44.64 q/ha) being at par with 15th November both were significantly more over 25th November and 5th December. Sowing of wheat at 5th November and 15th

November produces 6.13%, 3.70% more grain yield over 25th November and 33.65% and 30.59% more grain yield over 5th December, respectively. Straw yield (52.90q/ha) recorded maximum with 5th November being at par with 15th November both were significantly more over 25th November and 5th December. Sowing of wheat at 5th November and 15th November produces 8.49%, 7.51% more straw yield over 25th November and 15.96% and 14.91% more straw yield over 5th December, respectively (Table 1). Similar findings was also reported by Dwivedi *et al.* (2015) [10].

The benefit accrued was more in wheat sowing at 5th November and statically at par with 15th November over 25th November and 5th December. Net return (Rs 39228/ha) recorded significantly higher with 5th November and statically at par with 15th November over 25th November and 5th December. The B-C ratio also recorded significantly higher with 5th November and statically at par with 15th November over 25th November and 5th December. The higher returns by 6.13 and 3.71% & by 33.65% & 30.60%, respectively were recorded when wheat sown at 5th November and 15th November than 25th November and 5th December (Table 2).

Table 1: Effect of date of sowing on yield and yield attributes and yield on wheat (Pooled data of two years)

Treatment	No. of effective tillers/m ²	Length of Spike(cm)	No of grain/spike	1000 grain Weight(g)	Grain yield (q/ha)	Straw yield (q/ha)
T ₁ - 5 November	334.20	9.54	48.40	44.56	44.64	52.90
T ₂ -15 November	318.20	9.13	44.80	45.06	43.62	52.42
T ₃ -25 November	276.40	8.79	43.80	43.80	42.06	48.76
T ₄ -5 December	235.00	8.25	40.20	41.50	33.40	45.62
LSD (P=0.05)	22.97	0.37	2.98	0.91	2.32	2.85

Table 2: Effect of date of sowing on economics on wheat (Pooled data of two years)

Treatment	Cost of cultivation(Rs/ha)	Gross return(Rs/ha)	Net return (Rs/ha)	B:C Ratio
T ₁ - 5 November	25500	64,728	39,228	2.54
T ₂ -15 November	25500	63,249	37,749	2.48
T ₃ -25 November	25500	60,987	35,487	2.39
T ₄ -5 December	25500	48,430	22,930	1.90
LSD (P=0.05)	-	3,367	3,367	0.13

References

1. IPCC (Intergovernmental Panel on Climate Change). Intergovernmental Panel on Climate Change fourth assessment report: Climate change 2007. Synthesis Report. World Meteorological Organization, Geneva, Switzerland; c2007.
2. Joshi AK, Chand R, Arun B, Singh RP and Ortiz Ferrara G. Breeding crops for reduced-tillage management in the intensive, rice-wheat systems of South Asia. *Euphytica*. 2007a;153:135-151.
3. Joshi AK, Mishra B, Chatrath R, Ortiz Ferrara G and Singh RP. Wheat improvement in India: present status, emerging challenges and future prospects. *Euphytica*. 2007b;157:431-446.
4. Kumar P, Sarangi A, Singh DK, Parihar SS. Wheat performance as influenced by saline irrigation regimes and cultivars. *Journal of Agri Search*. 2014;1(2):66-72.
5. Meena BL, Singh AK, Phogat BS, Sharma HB. Effects of nutrient management and planting systems on root phenology and grain yield of wheat. *Indian J. Agril. Sci*. 2013;83(6):627-632.
6. Mian MA, Mahmood A, Ihsan M, Cheema NM. Response of different wheat genotypes to post anthesis temperature stress. *J Agric. Res*. 2007;45:269-276.
7. Singh AK, Manibhushan, Chandra N, Bharati RC. Suitable crop varieties for limited irrigated conditions in different agro climatic zones of India. *Int. J Trop Agr*. 2008;26(3-4):491-496.
8. Srivastava N, Singh D, Shukla A, Guru SK, Singh M, Rana DS. Effect of high temperature stress at post anthesis stage on photo system II, senescence, Yield attributes of wheat genotypes. *Indian J. Plant Physiol*. 2012;17:158-165.
9. Wardlaw IF, Blumenthal C, Larroque O, Wrigley CW. Contrasting effects of chronic heat stress and heat shock on grain weight and flour quality in wheat. *Functional Plant Biol*. 2002;29:25-34.
10. Dwivedi SK, Kumar Santoshand Prakash Ved. Effect of late sowing on yield and yield attributes of wheat genotypes in Eastern Indo Gangetic Plains (EGIP), *Journal of Agri Search*. 2015;2(4):304-306.