



ISSN (E): 2277-7695  
ISSN (P): 2349-8242  
NAAS Rating: 5.23  
TPI 2023; 12(7): 2121-2122  
© 2023 TPI  
[www.thepharmajournal.com](http://www.thepharmajournal.com)  
Received: 28-04-2023  
Accepted: 30-05-2023

**Sonia Dansena**  
Department of Fruit Science,  
Pt K.L.S. CHRS, IGKV, Raipur,  
Chhattisgarh, India

**Dr. Shishir Prakash Sharma,**  
Assistant Professor, Department  
of Fruit Science, Pt K.L.S.  
CHRS, IGKV, Raipur,  
Chhattisgarh, India

## To study about biochemical changes (pH, ascorbic acid and Total sugar %) of dragon fruit nectar during storage

**Sonia Dansena and Dr. Shishir Prakash Sharma**

### Abstract

The current investigatory work entitled "study about biochemical changes (pH, ascorbic acid and Total sugar %) of dragon fruit nectar during storage" was executed at Pomology Laboratory, Department of Pomology, Pt K.L.S. CHRS, Pendri, Rajnandgaon, IGKV, Raipur, Chhattisgarh during year 2019-2020. The analysis of chemical variation of nectar of dragon fruit at fortnight meantime for course of 45 days preservation by implementing CRD with 3 replication inside lab condition. Each replication comprised of 7 treatments for nectar. The nectar comprise of 20 percent pulp and 0.3 percent acidity and recipe is varied by different concentration of TSS. The nectar prepared from treatment T<sub>7</sub> containing 20 percent pulp, 0.3 percent acidity and 22 percent TSS shows highest value of ascorbic acid, TSS, total sugar, while The pH value of Nectar was peak for treatment T<sub>4</sub> comprising of 20% pulp, 0.3% acidity, 19% TSS.

**Keywords:** Dragon fruit (*Hylocereus costaricensis*), pH, ascorbic acid, total sugar, nectar, replication, treatment, biochemical

### 1. Introduction

Dragon fruit (*Hylocereus* spp.) is a sweltering climate bearer climbing cactus. The genus *Hylocereus* belongs to Cactaceae family, which is a dicotyledonous flowering plant family, under Caryophyllales order. In Latin America it is also known belle of the night and condrella plant. It contains antioxidant such as flavanoids, phenolic acid and betacyanin and naturally fat free and high fibre. These is a low acid food and Its pH values ranges between 4.4 and 5.1 out of which malic acid forms major portion. (Nomura *et al.*, 2005) [2]. Biochemical estimation of fruit showed that the 100 gm fruit has moisture content of about 83-88%, titratable acidity between 0.20 to 0.30 mg lactic acid equivalents, total soluble solids (TSS) between 8-12°Brix. Dragon fruit can be consumed fresh. Its flowers can be consumed directly or macerated in tea. Rarely utilized in flavouring other drinks. The pulpy and luscious flesh can also be jumbled with milk or sugar, utilized in marmalades, jellies, ices and soft drinks. It can also be transformed into juice, jam, RTS, nectar, squash, red wine, etc. Beside it can be useful to develop factory products such as preserve, ice cream, syrup, yogurt, candy, pastry, spread and ketch up. (Anonymous, 2006) [1]. Rarely pulp jumbled in pizzas. Processed products can be made from fresh pulp or frozen pulp. The red and pink pulp of dragon fruit can be used as a food dyeing agent and as a raw material for the food dyeing factories (Wybraniec and Mizrahi, 2002) [4]. The flower buds are utilized to make soups or jumbled in salads, and could also be consumed as a vegetable. The peels can be parched to separate pectin and antioxidants. The utilization of red pitaya as a raw material makes the food attractive and eye catching and also add up to nutritional composition of its processed products.

This fruit is new to Chhattisgarh and is cultivated in some parts of Raigarh district (kotra, Kharsia) Raipur, Rajnandgaon and Durg district. And there are rare work done in dragon fruit in India especially in processing. As hot climate and low rainfall is suitable for its cultivation so its area and production will increase in Chhattisgarh in future so there is need to standardize the recipe for preparation of useful products from it such as jam, jelly and beverages.

### 2. Materials and Method

The recent research on study about biochemical changes (pH, ascorbic acid and Total sugar %) of dragon fruit nectar during storage was conducted during 2019-20 in the laboratory of Department of fruit science, Pt K.L.S College of Horticulture and Research Station Rajnandgaon (C.G).

**Corresponding Author:**  
**Sonia Dansena**  
Department of Fruit Science,  
Pt K.L.S. CHRS, IGKV, Raipur,  
Chhattisgarh, India

It is located at 21.10°N latitude, and 81.03°E longitude and at an altitude of 307 m MSL under Chhattisgarh plains. It has tropical and dry spell throughout the year, However the temperature observed is 10 °C during winter and in summer reaches the 48 °C. This place bears hot windy climate during summer and precipitation annually of 1250 mm out of which approximately 85% is precipitated from third week of June to mid of September and remaining precipitate in month of October to February. May month has maximum temperature while December month has minimum temperature. The basic material used in this present research are firm, well developed and uniform ripened dragon fruit of *Hylocerus costaricensis* species which were obtained from farm of Chawda Bagh, Nandanvan Road, Raipur. And chemical and solution needed for analysis were provided by Fruit Science Processing laboratory, Pt. K.L.S College of Horticulture and Research Station Rajnandgaon C.G and Soil Science laboratory, S.K.S College of Agriculture and Research Station, Rajnandgaon C.G. Equipment and instruments required for analysis of product are provided by Soil Science laboratory, S.K.S College of Agriculture and Research Station, Rajnandgaon C.G.

The bottles of nectar beverage was stored under favourable environment for further analysis and observation upto 45 days and analysis conducted at 15 days interval. The pH was evaluated by usage a digital pH meter after callibrating it with buffer pH 4.0 and 9.2, ascorbic acid content of Nectar are analysed by the standard methods described by Ranganna (1997) [3], sugar was estimated by procedure of Lane and Eynon as described by Ranganna (1997) [3].

### 3. Result and Discussion

The dragon fruit nectar shows following biochemical variation during storage At its intial preparation time peak pH value observed was 4.72 with the treatment T<sub>4</sub> (20 percent pulp+19 percent TSS + 0.3 percent acidity), After 15 days storage of dragon fruit Nectar peak pH value observed was 4.63 with the treatment T<sub>4</sub> (20 percent pulp+13 percent TSS + 0.3 percent acidity), After 30 days storage of dragon fruit Nectar peak pH value observed was 4.48 with the treatment T<sub>4</sub> (20 percent pulp +13 percent TSS + 0.3 percent acidity), After 45 days storage of dragon fruit Nectar peak pH value observed was 4.32 with the treatment T<sub>4</sub> (20 percent pulp+13 percent TSS + 0.3 percent acidity).

At its intial preparation time peak ascorbic acid value observed was 11.30 with the treatment T<sub>7</sub> (20% pulp+22 % TSS + 0.3% acidity), after 15 days storage of dragon fruit Nectar peak ascorbic acid value observed was 10.03 with the treatment T<sub>7</sub> (20 percent pulp+ 22 percent TSS + 0.3 percent acidity), after 30 days storage of dragon fruit peak ascorbic acid value observed was 5.90 with the treatment T<sub>7</sub> (20 percent pulp + 22 percent TSS + 0.3 percent acidity), after 45 days storage of dragon fruit Nectar peak ascorbic acid value observed was 4.70 with the treatment T<sub>7</sub> (20 percent pulp + 22 percent TSS + 0.3 percent acidity).

At its intial preparation time peak total sugar value observed was 12.31 with the treatment T<sub>7</sub> (20 percent pulp + 22 percent TSS + 0.3 percent acidity),after 15 days storage of dragon fruit Nectar peak total sugar value observed was 12.33 with the treatment T<sub>7</sub> (20 percent pulp+22 percent TSS + 0.3 percent acidity),after 30 days storage of dragon fruit Nectar peak total sugar value observed was 12.35 with the treatment T<sub>7</sub> (20 percent pulp+22 percent TSS + 0.3 percent

acidity),after 45 days storage of dragon fruit Nectar peak total sugar value observed was 12.37 with the treatment T<sub>7</sub> (20 percent pulp+ 22 percent TSS + 0.3 percent acidity).

### 4. Conclusion

It is evident from findings that the pH and ascorbic acid value of Nectar shows diminishing trend with enhancement in time of storage (0-45 days), while the total sugar value of Nectar shows growing pattern with enhancement in course of preservation (0-45 days).

### 5. References

1. Anonymus. Department of agriculture: Crop Recommendations. Retrieved September 22, 2014, from Department of Agriculture; Government of Sri Lanka; c2006.
2. Nomura K, Idle M, Yonemoto Y. Changes in sugars and acids in pitaya (*Hylocerus. undatus*) fruit during development, The journal of Hort. Sci. and Biotech. 2005;80(6):711-715.
3. Ranganna S. Handbook of Analaysis and Quality Control for Fruit and Vegetable Products. Tata McGraw Hill, New Delhi; c1977.
4. Wybraniec S, Mizrahi Y. Fruit flesh betacyanin pigments in *Hylocereus cacti*. Journal of Agricultural and Food Chemistry. 2002;50:6086-6089.